

# Fractional Integration Operators in Mixed Weighted Generalized Hölder Spaces of Function of Two Variables Defined By Mixed Modulus of Continuity

T. Mamatov

Department of Higher mathematics, Bukhara Technological Institute of Engineering, Bukhara, Uzbekistan

**Corresponding Author:** T. Mamatov, Department of Higher mathematics, Bukhara Technological Institute of Engineering, Bukhara, Uzbekistan, E-Mail: [mamatov.tulkin@mail.ru](mailto:mamatov.tulkin@mail.ru)**Received date: march 16, 2020; Accepted date: April 23, 2020; Published date: May 29, 2020****Citation:** T. Mamatov (2020) Fractional Integration Operators in Mixed Weighted Generalized Hölder Spaces of Function of Two Variables Defined by Mixed Modulus of Continuity. J Mathematical Methods in Engineering. 1(1): DOI:[10.31579/jmme.2019/004](https://doi.org/10.31579/jmme.2019/004)**Copyright:** © 2020 T. Mamatov. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Abstract

In the presented work for operators the mixed fractional integration character of improvement of smoothness in comparison with smoothness of density  $\varphi(x, y)$  with weight  $\rho(x, y)$  in case of its any continuity modulus is found out  $\omega(\rho\varphi; x, y)$ . Zygmund type estimates are received. We consider operators of mixed fractional integration in weighted generalized Hölder spaces of a function of two variables defined by a mixed modulus of continuity.

**Keywords:** function of two variable, mixed fractional integral, mixed difference, generalized holder space, weighted space, weight, and mixed modulus of continuity

## 1. Introduction

One of the most important problems in the theory of integral operators in space is the problem of elucidating the dependence of the smoothness of the image on the smoothness of the preimage. The solution to such a problem plays an important role in the solvability of integral equations, their stability, and so on. The concept of smoothness can be formulated in a variety of terms. One of the ways of sufficiently fine-grabbing the smoothness of functions is the notion of generalized Hölderness, formulated in terms of the behavior of the modulus of continuity. Thus, one of the important questions in the theory of operators is as follows: Let be  $A$  an operator acting in a Banach space  $X$  and let be the modulus of continuity  $\omega(f; h) = \sup_{|t| \leq h} |f(x+t) - f(x)|_X$  of  $X$ . How can the

behavior of the modulus of continuity be characterized if the behavior of the modulus of continuity of function  $\omega(\varphi; h): \omega(\varphi; h) \leq C\psi(h)$  for all is known  $\varphi \in X$ , where  $\psi(x)$  a given continuous function,  $\psi(0) = 0$  the weight case. This problem admits a natural generalization to the weight case namely, let  $\rho(x)$  - weight function and  $\omega(\rho\varphi, h) \leq C\psi(h)$  for all  $\varphi \in X$ . How to estimate the modulus of continuity  $\omega(\rho A\varphi, h)$ ? A similar problem can be considered completely solved for

different spaces, and also for the Holder space of functions of one variable  $(A^\alpha \varphi)(x) = \frac{1}{\Gamma(\alpha)} \int_0^x \varphi(t) dt^{\alpha}$ ,  $0 < \alpha < 1$  ([2] – [6], [13] – [19]).

A detailed review of these and some other close results can be found in [12].

The assertion for multidimensional cases on the property of mapping in the usual Hölder and in the Hölder spaces defined by mixed differences are known [7], [8], [9], [10], [11], [12].

A similar problem in generalized Hölder spaces of the function of several variables has not been studied. This paper is aimed to fill in this gap. We deal with both non-weighted and weighted spaces. An important stage in the study of fractional integro-differentiation of functions from generalized Hölder spaces (see [1] - [6], [13], [14], [18]) is obtaining estimates of Zygmund type; Estimate of the modulus of continuity of a fractional integral in terms of the modulus of continuity of the original function. The main thrust of the work is to obtain an estimate of the Zygmund type that majorizes the mixed modulus  $\omega(\rho I_{+,+}^{\alpha,\beta} \varphi; h, \eta)$  of continuity of a mixed fractional integral with the weight of integral constructions from the mixed modulus of continuity  $\omega(\rho\varphi; h, \eta)$  of its density  $\varphi(x, y)$  with weight  $\rho(x, y)$ . These Zygmund-type estimates and action theorems directly affect the character of the improvement of the modulus of continuity by a mixed fractional integration  $I_{+,+}^{\alpha,\beta} \varphi$  of order  $(\alpha, \beta)$ .



$$(I_{0+,0+}^{\alpha,\beta}\varphi)(x,y) = \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_0^x \int_0^y \frac{\varphi(t,\tau)dtd\tau}{(x-t)^{1-\alpha}(y-\tau)^{1-\beta}}, \quad x>0, y>0 \quad (1)$$

where  $x>0, y>0$  and  $\alpha, \beta \in (0,1)$ .

It should be emphasized that the presence of weight significantly affects the nature of the Zygmund type evaluation. This was known in the case of Zygmund type estimates for fractional integrals of functions of one variable.

This paper is devoted to the study of certain properties of the mixed fractional integral (1) in weighed generalized Hölder spaces of a function of two variables defined by mixed modulus of continuity.

We consider the operator (1) in  $Q = \{(x,y) : 0 < x < b, 0 < y < d\}$ .

## 2. Preliminary information and notations

When studying the properties of continuous functions of several variables, in particular, two variables, the following classes of functions arise:

$$H^{\omega_1, \omega_2, \omega_{1,1}} = \left\{ \begin{array}{l} \varphi(x,y) \in C \\ \omega(\varphi; \delta, 0) = O(\omega(\delta)), \omega(\varphi; \sigma) = O(\omega(\sigma)), \\ \omega(\varphi; \delta, \sigma) = O(\omega_{1,1}(\delta, \sigma)) \end{array} \right\}_Q^{1,0} \quad (1)$$

$$H^{\omega_1, \omega_2} = \left\{ \begin{array}{l} \varphi(x,y) \in C \\ \omega(\varphi; \delta, 0) = O(\omega(\delta)), \omega(\varphi; \sigma) = O(\omega(\sigma)), \\ \omega(\varphi; \delta, \sigma) = O(\omega_{1,1}(\delta, \sigma)) \end{array} \right\}_Q^{1,0} \quad (2)$$

Where  $\omega(\varphi; \delta, 0) = \sup_{y \in [0, \delta]} \sup_{h \in (0, \delta]} \left| \Delta_h^{\omega_1} \varphi(x, y) \right|$ ,  $\omega(\varphi; 0, \sigma) = \sup_{x \in [0, \sigma]} \sup_{\eta \in (0, \sigma]} \left| \Delta_\eta^{\omega_2} \varphi(x, y) \right|$  - are the partial modulus of continuity of the first order, and  $\omega(\varphi; \delta, \sigma) = \sup_{x \in [0, \sigma]} \sup_{\eta \in (0, \sigma]} \left| \Delta_{h, \eta}^{\omega_{1,1}} \varphi(x, y) \right|$  is mixed modulus of continuity of order

$$\left( \Delta_h^{\omega_1} \varphi \right)(x, y) = \varphi(x+h, y) - \varphi(x, y), \quad \left( \Delta_\eta^{\omega_2} \varphi \right)(x, y) = \varphi(x, y+\eta) - \varphi(x, y),$$

$$\left( \Delta_{h, \eta}^{\omega_{1,1}} \varphi \right)(x, y) = \varphi(x+h, y+\eta) - \varphi(x+h, y) - \varphi(x, y+\eta) + \varphi(x, y),$$

$\omega_1, \omega_2 \in \Phi^1$ ,  $\omega_{1,1} \in \Phi^{1,1}$  (Definition of classes  $\Phi^1$  and  $\Phi^{1,1}$  see below).

$$\varphi(x+h, y+\eta) = \left( \Delta_{h, \eta}^{\omega_{1,1}} \varphi \right)(x, y) + \left( \Delta_h^{\omega_1} \varphi \right)(x, y) + \left( \Delta_\eta^{\omega_2} \varphi \right)(x, y) + \varphi(x, y). \quad (2)$$

**Definition 1.** Let function  $\varphi(x)$  is a bounded on  $[a, b]$ . The modulus of continuity of  $\varphi(x)$  is the expression

$$\omega(\varphi; \delta) = \sup_{\substack{x_1, x_2 \in [a, b] \\ |x_1 - x_2| \leq \delta}} |\varphi(x_1) - \varphi(x_2)|$$

is defined for all  $\delta$  that satisfy the condition  $0 < \delta \leq b - a$ .

**Definition 2.** A function  $\omega(\delta)$  is called a modulus of continuity if it satisfies conditions

- 1)  $\lim_{\delta \rightarrow 0} \omega(\delta) = 0$ ;
- 2)  $\omega(\delta)$  is almost increasing on  $(0, b-a]$ ;
- 3)  $\omega(\delta_1 + \delta_2) \leq \omega(\delta_1) + \omega(\delta_2)$ ;
- 4)  $\omega(\delta)$  is function continuous in  $\delta$  on  $(0, b-a]$ .



**Definition 3.** We denote by  $\Phi^1$  the class of functions  $\omega(\delta)$  defined on  $(0, b-a]$  and satisfying conditions

- a)  $\omega(\delta) \sim_{\underline{\omega}(\delta)}$  modulus of continuity
- b)  $\int_{\delta}^{b-a} \underline{\delta}(t) dt \leq C\omega(\delta);$
- c)  $\delta \int_{\delta}^{b-a} \frac{dt}{t^2} \leq C\omega(\delta);$
- d)  $\omega'(\delta) \sim_{\delta} .$

It follows from the definition  $\omega^{1,1}(\varphi; \delta, \sigma)$  that this function belongs to  $\Phi^1(Q)$  each of the variables. In addition, we note

$$\omega^{1,1}(\varphi; \delta, \sigma) \leq 2 \min \left\{ \omega^{1,0}(\varphi; \delta, 0), \omega^{0,1}(\varphi; 0, \sigma) \right\}. \quad (3)$$

**Definition 4.** We denote by  $\Phi^{1,1}(Q)$  the class of functions of two variables  $\omega(\delta, \sigma)$  satisfying conditions:

- 1)  $\omega(\delta, \sigma)$  in  $\delta$  for any fixed  $\sigma$ ;
- 2)  $\omega(\delta, \sigma)$  in  $\sigma$  for any fixed  $\delta$ .

We call this class the class of mixed modulus of continuity of the first order of continuous functions of two variables.

In [1] was shown that the properties 1) and 2) are characteristic for continuity modulus in the sense that for every

$$\omega \in \Phi^{1,1}(Q) \text{ there exist such a function } \varphi \in C_Q, \text{ that}$$

$$\omega^{1,1}(\varphi; \delta, \sigma) \sim \omega^{1,0}(\delta, \sigma), \omega^{1,0}(\varphi; \delta, 0) \sim \omega^{0,1}(\delta), \omega^{0,1}(\varphi; 0, \sigma) \sim \omega^{0,1}(\sigma).$$

**Definition 5.** Let us denote  $\Phi(Q)$  the set of satisfying  $(\omega_1, \omega_2, \omega_{1,1})$

- 1)  $\omega_1(\delta), \omega_2(\sigma) \in \Phi^1;$
- 2)  $\omega_{1,1}(\delta, \sigma) \in \Phi^{1,1};$
- 3)  $\omega_{1,1}(\delta, \sigma) \leq C \min \left\{ \omega_1(\delta), \omega_2(\sigma) \right\},$

Where  $C$  – is not envy from  $\omega_1, \omega_2, \omega_{1,1}$ .

Let  $\omega = (\omega_1, \omega_2, \omega_{1,1}) \in \Phi = \Phi^1 \times \Phi^1 \times \Phi^{1,1}$ . We have introduced a norm in  $\tilde{H}^\omega = H^{(\omega_1, \omega_2, \omega_{1,1})}$  space

$$\|\varphi\|_{\tilde{H}^\omega} = \max \left\{ \|\varphi\|_{C(Q)}, C_{\omega_1}, C_{\omega_2}, C_{\omega_{1,1}} \right\},$$

Where

$$C^{1,0} = \sup_{\varphi} \frac{\omega_1(\varphi; \delta, 0)}{\omega_1(\delta)}, \quad C^{0,1} = \sup_{\varphi} \frac{\omega_2(\varphi; 0, \sigma)}{\omega_2(\sigma)}, \quad C^{1,1} = \sup_{\varphi} \frac{\omega_{1,1}(\varphi; \delta, \sigma)}{\omega_{1,1}(\delta, \sigma)},$$

$$\|\varphi\|_{\tilde{H}^\omega} = \max_{(x,y) \in Q} |\varphi(x, y)|.$$

**Definition 6.** We say that  $\varphi(x, y) \in H^\omega(Q)$ , if  $\varphi(x, y) \in H^1(Q)$  and  $\varphi(x, y)|_{x=0, y=0} = \varphi(x, y)|_{x=b, y=d} = 0$ .

We will also make use of the following weighted spaces. Let  $\rho(x, y)$  be a non-negative function on  $Q$  (we will only deal with degenerate weights  $\rho(x, y) = \rho(x)\rho(y)$ ).

**Definition 7.** By  $\tilde{H}^\omega(Q, \rho) = \tilde{H}^\omega(\rho)$  we denote the space of functions  $\varphi(x, y)$  such that  $\rho\varphi \in \tilde{H}^\omega$ , respectively, equipped with the norm



$$\|\varphi\|_{H^{\omega}(\rho)} = \|\rho\varphi\|_H.$$

By  $\tilde{H}_0^{\omega}(\rho)$  we denote the corresponding subspaces of functions  $\varphi(x, y)$  such that

$$\rho(x, y)\varphi(x, y)|_{x=0, y=0} = \rho(x, y)\varphi(x, y)|_{x=b, y=d} = 0.$$

Below we follow some technical estimations suggested in [3] for the case of one-dimensional Riemann - Liouville fractional integrals. We denote

$$B(x, y; t, \tau) = \frac{\rho(x, y) - \rho(t, \tau)}{\rho(t, \tau)(x-t)^{1-\alpha}(y-\tau)^{1-\beta}},$$

Where  $0 < \alpha, \beta < 1$ ,  $0 < t < x < b$ ,  $0 < \tau < y < d$ . In the case  $\rho(x, y) = \rho(x)\rho(y)$  we have

$$B(x, y; t, \tau) = B_1(x, t)B_2(y, \tau) + \frac{B_1(x, t)}{(y-\tau)^{1-\beta}} + \frac{B_2(y, \tau)}{(x-t)^{1-\alpha}}, \quad (4)$$

Where

$$B_1(x, t) = \frac{\rho(x) - \rho(t)}{\rho(t)(x-t)^{1-\alpha}}, \quad B_2(y, \tau) = \frac{\rho(y) - \rho(\tau)}{\rho(\tau)(y-\tau)^{1-\beta}}.$$

Let also

$$D_1(x, h, t) = B_1(x+h, t) - B_1(x, t), \quad t, x, x+h \in [0, b], \quad h > 0$$

$$D_2(y, \eta, \tau) = B_2(y+\eta, \tau) - B_2(y, \tau), \quad \tau, y, y+\eta \in [0, d], \quad \eta > 0.$$

**Lemma 1.** ([3]) Let  $\rho(x) = x^\mu$ ,  $\mu \in \mathbb{R}^+$ ,  $0 < \alpha < 1$ . Then

$$|B_1(x, t)| \leq C \left( \frac{x}{t} \right)^{\max(\mu-1, 0)} \frac{(x-t)^\alpha}{t}, \quad (5)$$

$$|D_1(x, h, t)| \leq C \left( \frac{x+h}{t} \right)^{\max(\mu-1, 0)} \frac{h}{(x+h-t)^{1-\alpha} t}. \quad (6)$$

Similar estimates hold for  $B_2(y, \tau)$  and  $D_2(y, \eta, \tau)$  with  $\rho(y) = y^\nu$ .

**Remark 1.** All the weighted estimations of fractional integrals in the sequel are based on inequalities (5)-(4). Note that the right - hand sides of these inequalities have the exponent  $\max(\mu-1, 0)$  which means that in the proof it suffices to consider only the case  $\mu \geq 1$ , evaluations of  $\mu < 1$  being the same as for  $\mu = 1$ .

The following statements are known, begin first proved in (see also [17], p. 197). However, here we give a sketch of the proof of this lemma, in order to compose the representation of lightness for the two-dimensional case. Consider the one-

$$(I^\alpha \varphi)(x) = \frac{\text{one-dimensional fractional Riemann-Liouville integral}}{\Gamma(\alpha) \int_0^x (x-t)^{1-\alpha} dt}, \quad x > 0, 0 < \alpha < 1, \quad (7)$$

**Theorem 1.** Let  $\varphi(x)$  be continuous on  $[0, b]$  and  $\varphi(0) = 0$ . For the fractional integral (7), the estimate

$$\omega(I^\alpha \varphi, h) \leq Ch^b \frac{\omega(\varphi, t)}{t^{2-\alpha}} dt \quad (8)$$

$$(I^\alpha \varphi)(x) = \frac{1}{\Gamma(\alpha)} \int_0^x \frac{\varphi(0)}{(x-t)^{1-\alpha}} dt \stackrel{\text{Is valid.}}{\underset{\text{Proof Representing } \varphi \text{ as } \varphi = \varphi_1 + \varphi_2}{=}} \int_0^x \frac{\varphi_1(0)}{(x-t)^{1-\alpha}} dt + \int_0^x \frac{\varphi_2(0)}{(x-t)^{1-\alpha}} dt = A_1(x) + A_2(x).$$

Let  $h > 0$ ;  $x, x+h \in [0, b]$ . We have

$$A(x+h) - A(x) = \frac{\varphi(x) - \varphi(0)}{h} [(x+h)^\alpha - x^\alpha] + \frac{1}{\Gamma(\alpha)} \int_0^h (\varphi(x+t) - \varphi(t)) dt +$$

$$+ \frac{1}{\Gamma(\alpha)} \int_x^h [\varphi(x-t) - \varphi(t)] [(h+t)^{\alpha-1} - t^{\alpha-1}] dt = \Delta_1 + \Delta_2 + \Delta_3.$$

We have:  $|\Delta_1| \leq C \omega(\varphi; x) |(x+h)^\alpha - x^\alpha|$ . In the case  $x \leq h$  we have  $\Delta_1 \leq Ch \omega(\varphi; h)$ . Let  $x \geq h$ . Then

$$|\Delta_1| \leq C \omega(\varphi; x) x^\alpha \left[ \left(1 + \frac{h}{x}\right)^\alpha - 1 \right] \leq C \frac{\omega(\varphi; x)}{x^{1-\alpha}} h. \quad (9)$$

$$Cx \omega(\varphi; x) \leq \omega(\varphi; x) \int_x^{\alpha-1} t^{\alpha-2} dt \leq \int_x^{\alpha-1} t^{2-\alpha} dt \leq \int_h^{\alpha-1} t^{2-\alpha} dt.$$

It follows from (9) that

$$|\Delta_1| \leq Ch \int_x^h \frac{\omega(\varphi, t)}{t^{2-\alpha}} dt.$$

Further

$$|\Delta_2| \leq \int_0^h \frac{\omega(\varphi, t)}{t^{2-\alpha}} dt = h^\alpha \int_0^h \frac{\omega(\varphi, h\xi)}{(1-\xi)^{\alpha-1}} d\xi \leq Ch^\alpha \omega(\varphi; h),$$

$$\text{With } C = \int_0^1 (1-\xi)^{\alpha-1} d\xi.$$

$$\begin{aligned} \text{To estimate } \Delta_3 \text{ we distinguish the cases: } & 1) \quad x \geq h \quad \text{In the first case} \\ \Delta_3 & \leq C \left[ \int_0^h \omega(\varphi, t) [t^{\alpha-1} - (h+t)^{\alpha-1}] dt + \int_h^{\alpha-1} \omega(\varphi, t) [t^{\alpha-1} - (h+t)^{\alpha-1}] dt \right] \leq \\ & \leq C \left[ h^\alpha \omega(\varphi; h) + h \int_1^h \frac{\omega(\varphi, t)}{t^{2-\alpha}} dt \right]. \end{aligned}$$

$$\text{Obviously in the second case } \Delta_3 \leq Ch \omega(\varphi; h).$$

Estimates for  $\Delta_1, \Delta_2, \Delta_3$  lead to (8) if we take into account the fact that  $h^\alpha \omega(\varphi; h)$  is dominated by the right-hand side of (8). The latter is easily obtained in view of the monotonicity of the function  $\omega(\varphi; t)$ .

To obtain estimates of the Zygmund type in the weighted case, we use the notation and the proof scheme from [2] and [6].

**Theorem 2.** Let  $\rho(x) = x^\mu$ ,  $0 \leq \mu < 2 - \alpha$ . If the function  $f(x)$ ,  $x \in [a, b]$  satisfies the condition:

- 1)  $\rho(x) f'(x) dt$  and  $\rho(x) f(x)|_{x=0} = 0$ ;
- 2) the integral

$$\omega(\rho I_{0+}^\alpha f, h) \leq C \left( h^\alpha \int_0^{\alpha-1} \frac{\omega(\rho f, t)}{t^{\alpha+\gamma-1}} dt + h \int_1^h \frac{\omega(\rho f, t)}{t^{2-\alpha}} dt \right). \quad (10)$$

$(\rho I^\alpha f)(x) = (I^\alpha g)(x) + (J^\alpha g)(x)$ ,  $(J^\alpha g)(x) = \frac{1}{\Gamma(\alpha)} \int_0^x B(x, t) g(t) dt$ .

$$0+ \quad 0+ \quad 0+ \quad \Gamma(\alpha) \int_0^x$$



Here the estimates for  $(I_\alpha^\mu g)(x)$  are solved in Theorem 1. Now consider the difference where  

$$(J_{0+}^\alpha g)(x+h) - (J_{0+}^\alpha g)(x) = F_1(x, h) + F_2(x, h),$$
  
 Where

$$F_1(x, h) = \int_x^{x+h} B_1(x+h, t)g(t)dt, \quad F_2(x, h) = \int_x^x D_1(x, h, t)g(t)dt.$$

Taking into account Remark 1, we consider only the case  $1 \leq \mu < 2 - \alpha$ . From (5) we have

$$|F_1| \leq C \int_x^{x+h} \left( \frac{x+h}{t} \right)^{\mu-1} \frac{g(t)}{t} dt \leq C \int_x^{x+h} \left( \frac{x+h}{t} \right)^{\mu-1} \frac{\omega(g, t)}{t} dt.$$

$$|F_1| \leq Ch \int_x^{x+h} \frac{\omega(g, t)}{t^\mu} dt.$$

Using the property of almost decreasing  $\frac{\omega(g, t)}{t}$ , we obtain

$$|F_1| \leq Ch^{\mu+\alpha-1} \int_x^{x+h} \frac{\omega(g, t-x)}{(t-x)^\mu} dt = Ch^{\mu+\alpha-1} \int_0^h \frac{\omega(g, t)}{t^\mu} dt.$$

$$|F_1| \leq Ch^\alpha (x+h)^{\mu-1} \int_{-h}^0 \frac{\omega(g, x-t)}{(x-t)^\mu} dt \stackrel{\text{If } x > h, \text{ then}}{\leq} Ch^\alpha x^{\mu-1} \int_0^h \frac{\omega(g, x+t)}{t^\mu} dt \leq Ch^\alpha h^\mu \int_0^h \frac{\omega(g, x+t)}{t^\mu} dt.$$

$$|F_1| \leq Ch^\alpha h^\mu \int_0^h \frac{\omega(g, x+t)}{t^\mu} dt.$$

Collecting the estimates  $|F_1|$  we obtain the inequality for  $0 \leq \mu < 2 - \alpha$

$$\int_0^1 \frac{\omega(g, t)}{t^\gamma} dt,$$

We pass to the estimate  $F_2$ . Using the estimate (6), we obtain

$$|F_2| \leq Ch \int_0^x \left( \frac{x+h}{t} \right)^{\mu-1} \frac{\omega(g, t)}{(x+h-t)^{1-\alpha} t} dt. \quad (11)$$

$$|F_2| \leq Ch^{\alpha+\mu-1} \int_0^x \frac{\omega(g, t)}{t^\mu} dt \leq Ch^{\alpha+\mu-1} \int_0^h \frac{\omega(g, t)}{t^\mu} dt.$$

If  $h < x$ , then, we represent the right-hand side of (11) as a sum of three terms:

$$|F_2| \leq Ch \left[ \int_0^h + \int_h^x + \int_x^h \left( \left| \frac{x+h}{t} \right| \right)^{\mu-1} \frac{\omega(g, t)}{(x+h-t)^{1-\alpha} t} dt \right] = F'_1 + F'_2 + F''_2.$$

For the term  $F'_2$  the relations are valid  $x+h \leq 2(x+h-t)$ , therefore

$$F'_2 \leq Ch \int_0^h \frac{\omega(g, t) dt}{t^\mu (x+h-t)^{2-\alpha-\mu}} \leq Ch^{\alpha+\mu-1} \int_0^h \frac{\omega(g, t)}{t^\mu} dt.$$



For the summand,  $F_2''$  we have  $2t \leq x+h$ , so  $1 \leq \mu < 2-\alpha$  we obtain the estimate

$$F_2'' \leq Ch \int_h^{\frac{1}{2}(x+h)} \frac{\omega(g;t)(x+h)^{\mu-1}}{t^{\mu} \left| \frac{x+h-2}{2} \right|} dt \leq Ch \int_h^b \frac{\omega(g;t)}{t^{2-\alpha}} dt.$$

We estimate the term  $F_2'''$ . Here  $t \geq x+h-t$  therefore  $\frac{t}{x+h-t} \leq C$ , it follows that

$$F_2''' \leq Ch \int_{\frac{1}{2}(x+h)}^{xa} \left( \frac{x+h}{t} \right)^{\mu-1} \frac{\omega(g;t) dt}{t(x+h-t)^{1-\alpha}} \leq Ch \int_{\frac{1}{2}(x+h)}^x \frac{\omega(g;x+h-t)}{(x+h-t)^{2-\alpha}} dt.$$

Because  $x+h \leq 2t$ . Having made the change  $\xi = x+h-t$  and going back to the variable  $t$ , we get

$$F_2''' \leq Ch \int_h^b \frac{\omega(g;t)}{t^{2-\alpha}} dt.$$

From the estimates  $F_2', F_2'', F_2'''$  follows when  $h < x$

$$|F| \leq C \left( h^{\mu+\alpha-1} \int_0^h \frac{\omega(g;t)}{t^\mu} dt + h \int_h^b \frac{\omega(g;t)}{t^{2-\alpha}} dt \right).$$

$$|F| \leq C \left( h^{\gamma+\alpha-1} \int_0^h \frac{\omega(g;t)}{t^\gamma} dt + h \int_h^b \frac{\omega(g;t)}{t^{2-\alpha}} dt \right), \quad \gamma = \max(1, \mu),$$

Which completes the proof.

### Zygmund type estimates for the mixed fractional integral

**Theorem 3.** Let  $\varphi \in C(Q)$  and  $\varphi(x, y)|_{x=0, y=0} = 0$ . Then for (1), we have estimates of the Zygmund type

$$\omega(f; h, \eta) \leq C_1 h \eta \int_h^b \int_\eta^b \frac{\omega(\varphi; t, \tau)}{t^{2-\alpha} \tau^{2-\beta}} dt d\tau. \quad (12)$$

$$\omega(f; h, 0) \leq C_2 h \int_h^b \frac{\omega(\varphi; t, 0)}{t^{2-\alpha}} dt, \quad \omega(f; 0, \eta) \leq C_3 \eta \int_\eta^b \frac{\omega(\varphi; 0, \tau)}{\tau^{2-\beta}} d\tau. \quad (13)$$

**Proof.** Using the identity (2), we represent the integral (1) in the form

$$(I_{0+,0+}^{\alpha,\beta} \varphi)(x, y) = \frac{\varphi(0,0)}{\Gamma(1+\alpha) \Gamma(1+\beta)} + \frac{x^\alpha}{\Gamma(1+\alpha)} \psi_1(y) + \frac{y^\beta}{\Gamma(1+\beta)} \psi_1(x) + \psi(x, y),$$

$$\psi_1(x) = \frac{1}{\Gamma(\alpha)} \int_0^x \frac{\varphi(t,0) - \varphi(0,0)}{(x-t)^{1-\alpha}} dt, \quad \psi_1(y) = \frac{1}{\Gamma(\beta)} \int_0^y \frac{\varphi(0,\tau) - \varphi(0,0)}{(y-\tau)^{1-\beta}} d\tau,$$

$$\psi(x, y) = \frac{1}{\Gamma(\alpha) \Gamma(\beta)} \int_0^x \int_0^y \frac{\left( \Delta_{t,\tau}^{1,1} \varphi \right)(0,0)}{(x-t)^{1-\alpha} (y-\tau)^{1-\beta}} dt d\tau.$$

Let  $h > 0$ ;  $x, x+h \in [0, b]$ . Consider the difference

$$\begin{aligned} & \left( \Delta_{1,0}^{1,0} f \right) (x, y) = \frac{(x+h)^\alpha - x^\alpha}{\Gamma(\alpha)} \int_0^y g(x, y-\tau) d\tau + \\ & + \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_0^y \int_0^y g(x-t, y-\tau) - g(x, y-\tau) \frac{\tau^{1-\beta}(t+h)^{1-\alpha}}{dt d\tau} dt d\tau + \\ & + - \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_0^x \int_0^y g(x-t, y-\tau) - g(x, y-\tau) \frac{(t+h)^{\alpha-1} - t^{\alpha-1}}{\tau^{1-\beta}(t+h)^{1-\alpha}} |dt d\tau|. \end{aligned}$$

$$\begin{aligned} & \left( \Delta_h f \right) (x, y) \leq C \left| \left( x+h \right)^\alpha - x^\alpha \right| \omega(\varphi; x, y-\tau) d\tau + \int_0^y \frac{1}{\tau^{1-\beta}} \int_{-h}^0 \int_0^y \frac{1}{(h+t)^{1-\alpha}} \tau^{1-\beta} dt d\tau + \\ & + \int_0^y \int_0^y \omega(\varphi; t, y-\tau) |(h+t)^{\alpha-1} - t^{\alpha-1}| \tau^{1-\beta} dt d\tau. \end{aligned}$$

$$\begin{aligned} & \left( \Delta_{1,0}^{1,0} f \right) (x, y) \leq C \left| \left( x+h \right)^\alpha - x^\alpha \right| \omega(\varphi; t, 0) + \int_0^y \omega(\varphi; t, 0) dt + \\ & + \int_0^y \omega(\varphi; t, 0) |(h+t)^{\alpha-1} - t^{\alpha-1}| dt. \end{aligned}$$

We make use of (3) and obtain

$$\left( \Delta_h f \right) (x, y) \leq C_2 h \int_h^y \frac{1}{t^{2-\alpha}} dt. \quad (14)$$

Similarly, we can obtain the estimate

$$\left( \Delta_\eta f \right) (x, y) \leq C_2 \eta \int_0^y \frac{1}{\tau^{2-\beta}} d\tau. \quad (15)$$

From (14) and (15) follows the inequalities (13).

Let  $h, \eta > 0$  and  $x, x+h \in [0, b]$ ;  $y, y+\eta \in [0, d]$ . Consider the difference

$$\begin{aligned} & \left( \Delta_{1,1}^{h,\eta} f \right) (x, y) = \left( \Delta_{1,1}^{h,\eta} \right) \left[ (x+h)^\alpha - x^\alpha \right] \left[ (y+\eta)^\beta - y^\beta \right] + \\ & := \left( g(x, y) \right) \left[ (x+h)^\alpha - x^\alpha \right] \left[ (y+\eta)^\beta - y^\beta \right] + \\ & + \left( y+\eta \right)^\beta - y^\beta \int_0^h g(x+t, y) - g(x, y) \frac{\Gamma(1+\alpha)\Gamma(1+\beta)}{\Gamma(\alpha)\Gamma(1+\beta)} dt + \left( x+h \right)^\alpha - x^\alpha \int_0^\eta g(x, y+\tau) - g(x, y) \frac{\Gamma(\alpha+1)\Gamma(\beta)}{\Gamma(\alpha+1)\Gamma(\beta)} \int_0^h \left( \eta - \tau \right)^{1-\beta} dt + \\ & + \left( y+\eta \right)^\beta - y^\beta \int_0^h \left[ g(x-t, y) - g(x, y) \right] \left[ \frac{\Gamma(\alpha)\Gamma(1+\beta)}{\Gamma(\alpha+1)\Gamma(\beta)} \left[ (\tau+\eta)^{\beta-1} - \tau^{\beta-1} \right] \right] dt + \\ & + \frac{1}{\Gamma(1+\alpha)\Gamma(\beta)} \int_0^y g(x, y) \left[ \left( \frac{1}{h} - \frac{1}{y} \right) - \left( \frac{1}{h} - \frac{1}{y+\eta} \right) \right] \left[ \left( \frac{1}{h} - \frac{1}{y} \right)^{\alpha-1} - \left( \frac{1}{h} - \frac{1}{y+\eta} \right)^{\alpha-1} \right] d\tau. \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_0^h \int_0^y \int_0^{1,1} \left( \Delta_{t,\tau}^{1,1} g \right)(x,y) dt d\tau + \\
& + \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_0^h \int_0^y \int_0^{1,1} \left( \Delta_{t,-\tau}^{1,1} g \right)(x,y) (\tau+\eta)^{\beta-1} - \tau^{\beta-1} dt d\tau + \\
& + \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_0^x \int_0^y \int_0^{1,1} \left( \Delta_{t,\tau}^{1,1} g \right)(x,y) [(h+t)^{\alpha-1} - t^{\alpha-1}] dt d\tau + \\
& + \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_0^x \int_0^y \int_0^{1,1} \left( \Delta_{t,-\tau}^{1,1} g \right)(x,y) [h^{\alpha-1} - t^{\alpha-1}] dt d\tau. \\
& \left| \left( \Delta_{h,\eta}^{1,1} f \right)(x,y) \right| \leq C \left| \omega(\varphi; x, y) (x+h)^\alpha - x^\alpha \right| \left| (y+\eta)^\beta - y^\beta \right| + \\
& + \left| (y+\eta)^\beta - y^\beta \right| \int_0^y \frac{1,1}{(h-t)^{1-\alpha}} dt + \left| (x+h)^\alpha - x^\alpha \right| \int_0^y \frac{1,1}{(\eta-t)^{1-\beta}} d\tau + \\
& + \left| (y+\eta)^\beta - y^\beta \right| \int_0^x \frac{1,1}{\omega(\varphi; t, y)} \left[ (t+h)^{\alpha-1} - t^{\alpha-1} \right] dt + \\
& + \left| (x+h)^\alpha - x^\alpha \right| \int_0^y \frac{1,1}{\omega(\varphi; x, \tau)} \left[ (\tau+\eta)^{\beta-1} - \tau^{\beta-1} \right] d\tau + \\
& + \frac{h-\eta}{\omega(\varphi; t, \tau)} \left| \int_0^h \int_0^y \frac{1,1}{(h-t)^{1-\alpha}} \left( \eta-\tau \right)^{1-\beta} dt d\tau + \int_0^h \int_0^y \frac{1,1}{(h-t)^{1-\alpha}} \left( \eta-\tau \right)^{1-\beta} \left( \tau+\eta \right)^{\beta-1} - \tau^{\beta-1} dt d\tau \right| + \\
& + \int_0^x \int_0^y \int_0^{1,1} \frac{1,1}{(\tau-\eta)^{1-\beta}} \left| (h+t)^{\alpha-1} - t^{\alpha-1} \right| dt d\tau + \\
& + \int_0^x \int_0^y \int_0^{1,1} \omega(\varphi; t, \tau) \left| (t+h)^{\alpha-1} - t^{\alpha-1} \right| \left| (\tau+\eta)^{\beta-1} - \tau^{\beta-1} \right| dt d\tau.
\end{aligned}$$

Each term of this inequality is estimated in the standard way, and one can obtain

$$\left| \left( \Delta_{h,\eta}^{1,1} f \right)(x,y) \right| \leq Ch \int_1^b \int_1^d \int_{h\eta}^{1,1} \frac{\omega(\varphi; t, \tau)}{t^{2-\alpha}\tau^{2-\beta}} dt d\tau.$$

From which inequality (11) follows.

**Theorem 4.** Let  $\rho(x, y) = x^\mu y^\nu$ ,  $0 \leq \mu < 2-\alpha$ ,  $0 \leq \nu < 2-\beta$ . If the function  $\varphi(x, y) \in Q$  satisfies the following conditions:

$$1) \varphi_0(x, y) = \rho(x, y) \varphi(x, y) \in C(Q) \text{ and } \varphi_0(x, y)|_{x=0, y=0} = 0;$$

$$2) \int_0^b \int_0^d \frac{\omega(\varphi_0; t, \tau)}{t^\gamma \tau^\lambda} dt d\tau \quad \text{The integral converges for } \gamma = \max\{1, \mu\}, \lambda = \max\{1, \nu\}. \text{ Then the following estimates}$$

of Zygmund type are valid



$$\omega(\rho f; h, 0) \leq C \left[ h^{\alpha+\gamma-1} \int_0^h \frac{1}{t^\gamma} dt + h \int_h^b \frac{\omega(\rho \varphi; t, d)}{t^{2-\alpha}} dt \right], \quad (16)$$

$$\begin{aligned} \omega(\rho f; 0, \eta) &\leq C \left[ \eta^{\beta+\lambda-1} \int_0^\eta \frac{1}{\tau^\lambda} dt + \eta \int_\eta^d \frac{\omega(\rho \varphi; b, \tau)}{\tau^{2-\beta}} d\tau \right], \\ \omega(\rho f; h, \eta) &\leq C \left[ h^{\alpha+\gamma-1} \eta^{\beta+\lambda-1} \int_0^h \int_0^\eta \frac{(\rho \varphi; t, \tau)}{t^\gamma \tau^\lambda} dt d\tau + h \eta^{\beta+\lambda-1} \int_h^b \int_0^1 \frac{\omega(\rho \varphi; t, \tau)}{t^{2-\alpha} \tau^\lambda} dt d\tau + \right. \end{aligned} \quad (17)$$

$$\left. + h^{\alpha+\gamma-1} \eta \int_h^b \int_0^1 \frac{(\rho \varphi; t, \tau)}{t^\gamma \tau^{2-\beta}} dt d\tau + h \eta \int_h^b \int_0^1 \frac{\omega(\rho \varphi; t, \tau)}{t^{2-\alpha} \tau^{2-\beta}} dt d\tau \right]. \quad (18)$$

**Proof.** By Remark 1, it suffices to deal with the case  $\mu, \nu \geq 1$ . Let  $\varphi \in H_0(\rho)$  so that  $\varphi(x, y) = \varphi(x, y)\rho(x, y)$ .

$$\text{where } \varphi(x, y) \in H_0 \text{ and } \varphi(x, y) \Big|_{x=y=0} = 0. \text{ For } G(x, y) := \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \int_0^x \int_0^y \frac{(x-t)^{1-\alpha}(y-\tau)^{1-\beta}}{(x-t)^{1-\alpha}(y-\tau)^{1-\beta}} \varphi(t, \tau) dt d\tau.$$

$$\begin{aligned} G(x, y) &:= \frac{1}{\Gamma(\alpha)\Gamma(\beta)} \left( \int_0^x \int_y^x \frac{\varphi(t, \tau) dt d\tau}{(x-t)^{1-\alpha}(y-\tau)^{1-\beta}} + \int_0^y \int_x^y \frac{\varphi(t, \tau) dt d\tau}{(x-t)^{1-\alpha}(y-\tau)^{1-\beta}} \right) \\ &= G_1(x, y) + G_2(x, y). \end{aligned}$$

Here the question of the estimation of the modulus of continuity for the first term is solved by us in Theorem 3.

To estimate the term  $G_2(x, y)$  we note that the weight being degenerate, we have

$\rho(x, y) - \rho(t, \tau) = (\rho(x) - \rho(t))(\rho(y) - \rho(\tau)) + \rho(\tau)(\rho(x) - \rho(t)) + \rho(t)(\rho(y) - \rho(\tau))$ , which leads to the following representation

$$\begin{aligned} G(x, y) &= \int_0^x \int_0^y B_1(x, t) B_2(y, \tau) \varphi_0(t, \tau) dt d\tau + \int_0^x \int_0^y \frac{B_1(x, t) \varphi_0(t, \tau)}{(y-\tau)^{1-\beta}} dt d\tau + \\ &\quad + \int_0^y \int_0^x \frac{B_2(y, \tau) \varphi_0(t, \tau)}{(x-t)^{1-\alpha}} dt d\tau, \end{aligned}$$

Where the notation (4) has been used.

For the difference  $\Delta_h G(x, y)$  with  $h > 0$  and  $x, x+h \in [0, b]$ , we have

$$\begin{aligned} \Delta_h G(x, y) &= \int_x^{x+h} \int_y^2 B(x+h, t) B(y, \tau) \varphi(t, \tau) dt d\tau + \int_x^{x+h} \int_0^y D(x, h, t) B(y, \tau) \varphi(t, \tau) dt d\tau + \\ &\quad + \int_x^{x+h} \int_0^1 B(x+h, t) \frac{\varphi(t, \tau) dt d\tau}{(y-\tau)^{1-\beta}} + \int_x^{x+h} \int_0^1 D(x, h, t) \frac{\varphi(t, \tau) dt d\tau}{(y-\tau)^{1-\beta}} + \int_x^{x+h} \int_0^y B(y, \tau) \frac{\varphi(t, \tau) dt d\tau}{(x+h-t)^{\alpha-1}} + \\ &\quad + \int_x^{x+h} \int_0^2 B(y, \tau) \left[ (x+h-t)^{\alpha-1} - (x-t)^{\alpha-1} \right] \varphi(t, \tau) dt d\tau. \end{aligned}$$



Since  $\varphi_0(x,0)=0$  then the inequality

$$\begin{aligned} & \left| G_2(x+h,y) - G_2(x,y) \right| \leq \int_x^{x+h} \int_0^y |B_1(x+h,t)| |B_2(y,\tau)| \omega(\varphi_0;t,\tau) dt d\tau + \\ & + \int_{-h}^0 \int_0^y |\omega(\varphi_0;t,\tau)| |B_2(y,\tau)| dt d\tau + \int_x^y \int_0^1 |D_1(x,h,t)| |B_2(y,\tau)| \omega(\varphi_0;t,\tau) dt d\tau + \\ & + \int_x^0 \int_1^y |B_2(x+h,t)| \frac{\omega(\varphi_0;t,\tau)}{\tau} dt d\tau + \int_0^y \int_0^1 |D_2(x,h,t)| \frac{\omega(\varphi_0;t,\tau)}{\tau} dt d\tau + \\ & + \int_x^0 \int_0^1 |B_2(x+h,t)| \frac{(y-\tau)^{1-\beta}}{\tau} dt d\tau + \int_0^y \int_0^1 |D_2(x,h,t)| \frac{(y-\tau)^{1-\beta}}{\tau} dt d\tau + \\ & + \int_0^y \int_0^1 \omega(\varphi_0;t,\tau) |(h+t)^{\alpha-1} - t^{\alpha-1}| |B_2(y,\tau)| dt d\tau. \end{aligned}$$

We estimate integral  $\int_0^y |B_2(y,\tau)| \omega(\varphi_0;t,\tau) dt d\tau$ . Let  $1 \leq v < 2 - \beta$ , then from (5) we have

$$\begin{aligned} & \int_0^y |B_2(y,\tau)| \omega(\varphi_0;t,\tau) dt d\tau \leq C \int_0^y \left( \frac{y}{\tau} \right)^{v-1} \frac{(y-\tau)^{\beta}}{\tau} \omega(\varphi_0;t,\tau) dt d\tau \leq \\ & \leq Cy^{\beta} \int_0^y \left( \frac{y}{\tau} \right)^{v-1} \frac{\omega(\varphi_0;t,\tau)}{\tau} dt d\tau \leq C \omega(\varphi_0;t,d). \end{aligned}$$

Now we estimate next integral

$$\int_0^y \frac{\omega(\varphi_0;t,\tau)}{\tau^{1-\beta}} d\tau \leq Cy^{\beta} \int_0^y \frac{\omega(\varphi_0;t,\tau)}{\tau} d\tau \leq C \omega(\varphi_0;t,d)$$

From here and the estimates  $\Delta_1, \Delta_2, \Delta_3$  of Theorem 1 and from the estimates  $F_1, F_2$  in Theorem 2, one can easily verify the

$$\left| \left( \Delta_h^{1,0} G \right)_{2,2}(x,y) \right| \leq C_1 \left[ h^{\alpha+\gamma-1} \int_0^h \frac{\omega(\rho\varphi;t,d)}{t^\gamma} dt + h^b \int_h^b \frac{\omega(\rho\varphi;t,d)}{t^{2-\alpha}} dt \right] \quad (19)$$

where  $\gamma = \max(1, \mu)$ .

$$\left| \left( \Delta_h^2 G \right)_{2,2}(x,y) \right| \leq C \left[ \eta^{\beta+\lambda-1} \int_0^{\eta} \frac{\omega(\rho\varphi;b,\tau)}{\tau^\lambda} d\tau + \eta^d \int_\eta^d \frac{\omega(\rho\varphi;b,\tau)}{\tau^{2-\beta}} d\tau \right], \quad (20)$$

The estimate  
Is symmetrically obtained, where  $\lambda = \max(1, v)$ .

For the mixed difference  $\left( \Delta_{h,\eta}^{1,1} G \right)_{2,2}(x,y)$  with  $h, \eta > 0$  and  $\forall x, x+h \in [a, b], \forall y, y+\eta \in [c, d]$

the appropriate representation leading to the separate evaluation in each variable without losses in another variable is as

$$\left( \Delta_{h,\eta}^{1,1} G \right)_{2,2}(x,y) = \int_x^{x+h} \int_y^{y+\eta} B_1(x) \left( \frac{y+\eta-h}{t} \right) B_2(y+\eta, \tau) \varphi(t, \tau) dt d\tau +$$

$$\begin{aligned}
& + \int_0^x \int_0^y D_1(x, h, t) D_2(y, \eta, \tau) \varphi_0(t, \tau) dt d\tau + \int_x^{x+h} \int_0^y B_1(x+h, t) D_2(y, \eta, \tau) \varphi_0(t, \tau) dt d\tau + \\
& + \int_0^x \int_0^y \int_0^1 D(x, h, t) B_2(y+\eta, \tau) \varphi_0(t, \tau) dt d\tau + \int_x^{x+h} \int_0^y \int_0^1 \frac{(B(x+h, t))^\beta}{(y+\eta-\tau)^{1-\beta}} \varphi_0(t, \tau) dt d\tau + \\
& + \int_0^x \int_0^y \int_0^1 B(x+h, t) \left[ (y+\eta-\tau)^{\beta-1} - (y-\tau)^{\beta-1} \right] \varphi_0(t, \tau) dt d\tau + \\
& + \int_x^{x+h} \int_0^y \int_0^1 D(x, h, t) (y+\eta-\tau)^{\beta-1} \varphi_0(t, \tau) dt d\tau + \\
& + \int_0^x \int_0^y \int_0^1 D(x, h, t) \left[ (y+\eta-\tau)^{\beta-1} - (y-\tau)^{\beta-1} \right] \varphi_0(t, \tau) dt d\tau + \\
& + \int_0^x \int_0^y \int_0^1 \int_{x+h}^y B_2(y+\eta, \tau) \left[ (x+h-t)^{\alpha-1} - (x-t)^{\alpha-1} \right] B_2(y+\eta, \tau) \varphi_0(t, \tau) dt d\tau + \\
& + \int_0^x \int_0^y \int_0^1 \int_{x+h}^y (x+h-t)^{\alpha-1} D_2(y, \eta, \tau) \varphi_0(t, \tau) dt d\tau + \\
& + \int_0^x \int_0^y \int_0^1 \int_{x+h}^y (x+h-t)^{\alpha-1} D_2(y, \eta, \tau) \varphi_0(t, \tau) dt d\tau + \\
& + \int_0^x \int_0^y \int_0^1 \int_{x+h}^y \left[ (x+h-t)^{\alpha-1} - (x-t)^{\alpha-1} \right] D_2(y, \eta, \tau) \varphi_0(t, \tau) dt d\tau. \\
& \left| \left( \Delta_{h, \eta}^{1,1} G \right)(x, y) \right| \leq C \int_x^{x+h} \int_0^y B_1(x+h, t) B_2(y+\eta, \tau) \omega(\varphi_0; t, \tau) dt d\tau + \\
& + \int_0^x \int_0^y D_1(x, h, t) D_2(y, \eta, \tau) \omega(\varphi_0; t, \tau) dt d\tau + \\
& + \int_x^{x+h} \int_0^y B_1(x+h, t) D_2(y, \eta, \tau) \omega(\varphi_0; t, \tau) dt d\tau + \\
& + \int_0^x \int_0^y \int_0^1 D_1(x, h, t) B_2(y+\eta, \tau) \omega(\varphi_0; t, \tau) dt d\tau + \\
& + \int_0^x \int_0^y \int_0^1 B_1(x+h, t) \left[ (\eta+\tau)^{\beta-1} - \tau^{\beta-1} \right] \omega(\varphi_0; t, \tau) dt d\tau + \\
& + \int_0^x \int_0^y \int_0^1 \int_{-\eta}^0 D_1(x, h, t) \frac{\omega(\varphi_0; t, \tau)}{(\eta+\tau)^{1-\beta}} dt d\tau + \int_0^x \int_0^y \int_0^1 B_2(y+\eta, \tau) \frac{\omega(\varphi_0; t, \tau)}{(h+t)^{1-\alpha}} dt d\tau + \\
& + \int_0^x \int_0^y \int_0^1 \int_{-\eta}^0 D_1(x, h, t) \left[ (\eta+\tau)^{\beta-1} - \tau^{\beta-1} \right] \omega(\varphi_0; t, \tau) dt d\tau +
\end{aligned}$$

$$\begin{aligned}
& + \int_0^y \int_0^y [(h+t)^{\alpha-1} - t^{\alpha-1}] B(y+\eta, \tau) \omega \varphi(t, \tau) dt d\tau + \\
& + \int_{-h}^0 \int_0^y D(y, \eta, \tau) \frac{\omega(\varphi; t, \tau)}{t^{\gamma}} dt d\tau + \int_0^2 \int_{x+h}^0 \frac{B(x+h, t)}{t^{\lambda}} \omega \varphi(t, \tau) dt d\tau + \\
& + \int_0^x \int_0^y [(h+t)^{1-\alpha} - t^{1-\alpha}] D(y, \eta, \tau) \frac{(\eta+\tau)^{1-\beta}}{t^{\lambda}} dt d\tau.
\end{aligned}$$

We omit the details of evaluation of each term in the above representation; it is standard via Lemma 1 and yields

$$\begin{aligned}
|\Delta_{h,\eta} G(x, y)| & \leq C \left[ h^{\alpha+\gamma-1} \eta^{\beta+\lambda-1} \int_0^h \int_0^{\eta} \frac{1}{t^{\gamma} \tau^{\lambda}} (\rho \varphi; t, \tau) dt d\tau + h \eta^{\beta+\lambda-1} \int_h^b \int_0^{\eta} \frac{1}{t^{2-\alpha} \tau^{\lambda}} (\rho \varphi; t, \tau) dt d\tau + \right. \\
& \left. + h^{\alpha+\gamma-1} \eta \int_0^h \int_0^{\eta} \frac{(\rho \varphi; t, \tau)}{t^{\gamma} \tau^{2-\beta}} dt d\tau + h \eta \int_h^b \int_0^{\eta} \frac{\omega(\rho \varphi; t, \tau)}{t^{2-\alpha} \tau^{2-\beta}} dt d\tau \right], \quad (21)
\end{aligned}$$

where  $\gamma = \max(1, \mu)$  and  $\lambda = \max(1, \nu)$ .

From the inequalities (21), (20), (19) and (12), (13) we obtain the corresponding estimates (16), (17) and (18).

### Mapping properties of the mixed fractional integration operators

in the space  $\tilde{H}_0^\omega(\cdot)$

In this section, we give a generalization of the theorem to the weighted.

**Theorem 5.** Let  $0 < \alpha, \beta < 1$ ,  $\rho(x, y) = x^\mu y^\nu$ ,  $0 \leq \mu < 2 - \alpha$  and  $0 \leq \nu < 2 - \beta$ . If  $\omega(x, y) \in \Phi(Q)$  and  $x^{\gamma} y^{\gamma} (x^{\gamma} y^{\gamma})^{\lambda} \omega(t, \tau) \in H_{1,1}$  assume that

$$1) \int_0^x \int_0^y \left| \frac{x^{\gamma} y^{\gamma} (x^{\gamma} y^{\gamma})^{\lambda}}{t^{\alpha} \tau^{\alpha} y^{\lambda}} \right| \frac{dt d\tau}{\omega(t, \tau)} \leq C_1 \omega(x, y), \quad (22)$$

$$2) \int_0^y \int_0^x \left| \frac{x^{\gamma} y^{\gamma} (x^{\gamma} y^{\gamma})^{\lambda}}{t^{\alpha} \tau^{\alpha} y^{\lambda}} \right| \frac{dt d\tau}{\omega(t, \tau)} \leq C_1 \omega(x, y), \quad (23)$$

where  $\gamma = \max(\mu-1, 0)$ ,  $\lambda = \max(\nu-1, 0)$ . Then the mixed fractional integral operator (1) is bounded from the

weight space  $\tilde{H}_0^\omega(\cdot)$  to the space  $\tilde{H}_0^{\omega_{\alpha, \beta}}(\rho)$  with the same weight and with the characteristic

$$\omega_{\alpha, \beta}(t, \tau) = t^\alpha \tau^\beta \omega(t, \tau). \quad \sim_{\omega_{\alpha, \beta}}$$

**Proof.** Let  $f = I_{0+, 0+} \varphi$  where  $\varphi \in H_0(\rho)$ . We will show that  $f \in H_0(\rho)$ . For this, it suffices to show that

$$\sup_{h>0} \frac{\omega(\rho f; h, 0)}{h^\alpha \omega_1(h)} = C_1 < \infty, \quad \sup_{\eta>0} \frac{\omega(\rho f; 0, \eta)}{\eta^\beta \omega_2(\eta)} = C_2 < \infty,$$

$$\sup_{h>0, \eta>0} \frac{\omega(\rho f; h, \eta)}{h^\alpha \eta^\beta \omega_{1,1}(h, \eta)} = C_3 < \infty.$$

From membership  $\omega(t, \tau)$  in the class  $\Phi(Q)$  and satisfaction of inequalities (22), (23) the convergence of the integrals follows



$$\int_0^b \frac{\omega(\rho\varphi; t, d)}{t^\gamma} dt, \quad \int_0^d \frac{\omega(\rho\varphi; b, \tau)}{\tau^\lambda} d\tau, \quad \int_0^b \int_0^d \frac{\omega(\rho\varphi; t, \tau)}{t^\gamma \tau^\lambda} dt d\tau.$$

Therefore, there are estimates of the Zygmund type from Theorem 4. Whence follows

$$\begin{aligned} \frac{(\rho f; h, 0)}{h^\alpha \omega_{(h, d)}} &\leq C_1 \left| \frac{h}{\omega_{(h, d)}} \int_0^{h^{-1}} \frac{\omega(\rho\varphi; t, d)}{t^\gamma} dt + \frac{h}{\omega_{(h, d)}} \int_{h^{-1}}^h \frac{\omega(\rho\varphi; t, d)}{t^\gamma} dt \right| \leq \\ &\leq C_1 \|\rho\varphi\|_{H_0^\alpha} \left\{ \frac{1}{\omega_{(h, d)}} \int_0^{h^{-1}} \frac{\omega(\rho\varphi; t, d)}{t^\gamma} dt + \frac{h}{\omega_{(h, d)}} \int_{h^{-1}}^h \frac{\omega(\rho\varphi; t, d)}{t^{2-\alpha}} dt \right\}. \\ \frac{(\rho f; 0, \eta)}{\eta^\beta \omega_{(b, \eta)}} &\leq C_2 \left| \frac{\eta}{\omega_{(b, \eta)}} \int_0^{h^{-1}} \frac{\omega(\rho\varphi; b, \tau)}{\tau^\lambda} d\tau + \frac{\eta}{\omega_{(b, \eta)}} \int_\eta^h \frac{\omega(\rho\varphi; b, \tau)}{\tau^{2-\beta}} d\tau \right| \leq \\ &\leq C_2 \|\rho\varphi\|_{H_0^\alpha} \left\{ \frac{\eta^{\lambda-1}}{\omega_{(b, \eta)}} \int_0^{h^{-1}} \frac{\omega(\rho\varphi; b, \tau)}{\tau^\lambda} d\tau + \frac{\eta^{1-\beta}}{\omega_{(b, \eta)}} \int_\eta^h \frac{\omega(\rho\varphi; b, \tau)}{\tau^{2-\beta}} d\tau \right\}. \\ \frac{\bar{\omega}(\rho f; h, \eta)}{h^\alpha \eta^\beta \omega_{(h, \eta)}} &\leq C_3 \left| \frac{h^{\gamma-1} \eta^{\lambda-1}}{\omega_{(h, \eta)}} \int_0^{h^{-1}} \int_0^h \frac{\omega(\rho\varphi; t, \tau)}{t^\gamma \tau^\lambda} dt d\tau + \right. \\ &\quad \left. + \frac{h^{\gamma-1} \eta^{1-\beta}}{\omega_{(h, \eta)}} \int_0^h \int_0^1 \frac{\omega(\rho\varphi; t, \tau)}{t^\gamma \tau^{2-\beta}} dt d\tau + \frac{h^{1-\alpha} \eta^{\lambda-1}}{\omega_{(h, \eta)}} \int_0^{h^{-1}} \int_0^h \frac{\omega(\rho\varphi; t, \tau)}{t^{2-\alpha} \tau^\lambda} dt d\tau + \right. \\ &\quad \left. + \frac{h^{1-\alpha} \eta^{1-\beta}}{\omega_{(h, \eta)}} \int_h^1 \int_0^1 \frac{\omega(\rho\varphi; t, \tau)}{t^{2-\alpha} \tau^{2-\beta}} dt d\tau \right| \leq C_3 \|\rho\varphi\|_{H_0^\alpha} \left\{ \frac{h^{\gamma-1} \eta^{\lambda-1}}{\omega_{(h, \eta)}} \int_0^{h^{-1}} \int_0^h \frac{\omega(t, \tau)}{t^\gamma \tau^\lambda} dt d\tau + \right. \\ &\quad \left. + \frac{h^{\gamma-1} \eta^{1-\beta}}{\omega_{(h, \eta)}} \int_0^{h^{-1}} \int_0^h \frac{\omega(t, \tau)}{t^\gamma \tau^{2-\beta}} dt d\tau + \frac{h^{1-\alpha} \eta^{\lambda-1}}{\omega_{(h, \eta)}} \int_0^{h^{-1}} \int_0^h \frac{\omega(t, \tau)}{t^{2-\alpha} \tau^\lambda} dt d\tau + \right. \\ &\quad \left. + \frac{h^{1-\alpha} \eta^{1-\beta}}{\omega_{(h, \eta)}} \int_0^{h^{-1}} \int_0^h \frac{\omega(t, \tau)}{t^{2-\alpha} \tau^{2-\beta}} dt d\tau \right\}. \end{aligned}$$

It's obvious that

$$\frac{\omega(\rho f; h, 0)}{h^\alpha \omega_{(h, d)}} \leq C_1 \|\rho\varphi\|_{H_0^\alpha}, \quad \frac{\omega(\rho f; 0, \eta)}{\eta^\beta \omega_{(b, \eta)}} \leq C_2 \|\rho\varphi\|_{H_0^\alpha},$$

$$\frac{\omega(\rho f; h, \eta)}{h^\alpha \eta^\beta \omega_{(h, \eta)}} \leq C_3 \|\rho\varphi\|_{H_0^\alpha}. \quad (24)$$

We estimate  $\|\rho f\|_{C(Q)}$ . We have



$$\begin{aligned}\rho(x, y)f(x, y) &= x^\mu y^\nu \int \int \int \frac{\varphi_0(x-t, y-\tau)dtd\tau}{t^{1-\alpha} \tau^{1-\beta}} = \\ &= x^\alpha y^\beta \int \int \frac{\varphi^0[x-x\xi, y-ys]d\xi ds}{\xi^{1-\alpha}(1-s)^\mu s^{1-\beta}(1-s)^\nu}.\end{aligned}$$

Since  $\varphi_0(x, y)|_{x=0, y=0} = 0$ , then  
 $|\varphi_0(x-x\xi, y-ys) - \varphi_0(0, y-ys)| \leq C_1 \omega(\varphi_0; 1-\xi)$ ,  
 $|\varphi_0(x-x\xi, y-ys) - \varphi_0(x-x\xi, 0)| \leq C_2 \omega(\varphi_0; 1-s)$ ,  
 $\left| \int \int \frac{\varphi^0(x, y)}{\Delta_{x(1-\xi), y(1-s)}} d\xi ds \right| \leq C \omega(\varphi^0; 1-\xi, 1-s).$

It follows that

$$|\rho(x, y)f(x, y)| \leq C \|\varphi\|_{H_0} \int \int \frac{\omega_{1,1}(t, \tau) dtd\tau}{t^\mu \tau^{1-\alpha} (1-t)^{\nu} (1-\tau)^{1-\beta}}.$$

Therefore

$$\|\rho f\|_{C(Q)} \leq C \|\varphi\|_{\tilde{H}_0^\alpha}(\rho). \quad (25)$$

From the inequalities (24) and (25) follows the assertion of the theorem.

## References

- Baba-Zade M.A (1979) On class of continuous functions of two variables (Russian), Uch. Zam.MV i SSO AZSSR, ser. fiz-mat nauk, No 3, p. 32-41
- H.G. Hardy and J.E. Littlewood (1928) Some properties of fractional integrals, I. Mah. Z. 27, No 4, 565-606.
- N.K. Karapetians and L.D. Shankishvili (1999) A short proof of Hardy-Littlewood-type theorem for fractional integrals in weighted Hölder spaces. Fract. Calc. Appl. Anal. 2, No.2, p. 177-192.
- N.K. Karapetians and L.D. Shankishvili (2003) Fractional integro-differentiation of the complex order in generalized Hölder spaces. Integral Transforms Spec. Funct. 13, No. 3, p. 199-209.
- N.K. Karapetyants, Kh. M. Murdaev and A. Ya. Yakubov (1990) The isomorphism realized by fractional integrals in generalized Hölder classes. Dokl. Akad. Nauk SSSR 314, No 2, 288-291.
- N.K. Karapetyants, Kh. M. Murdaev and A. Ya. Yakubov (1992) On isomorphism provided by fractional integrals in generalized Nikolskiy classes. Izv. Vuzov. Matematika (9), 49-58.
- T. Mamatov, S. Samko (2010) Mixed Fractional Integration Operators in mixed weighted Hölder spaces. Fractional Calculus and Applied Analysis (FCAA), Vol. 13, No 3, pp. 245-259
- Mamatov T (2018) Mixed Fractional Integro-Differentiation Operators in Hölder Spaces. The latest research in modern science: experience, traditions and innovations. Proceedings of the VII International Scientific Conference. Section I. North Charleston, SC, USA. 20-21 June, P. 6-9
- Mamatov T (2014) Mapping Properties of Mixed Fractional Integro-Differentiation in Hölder Spaces. Journal of Concrete and Applicable Mathematics (JCAAM). Volume 12, Num. 3-4. p. 272-290
- Mamatov T., Rayimov D (2019) Elmurodov M. Mixed Fractioanl Differentiation Operators in Hölder Spaces. Journal of Multidisciplinary Engineering Science and Technology (JMEST), Vol. 6 Issue 4, P. 9855-9857
- Mamatov T., Homidov F., Rayimov D (2019) On Isomorphism Implemented By Mixed Fractional Integrals in Hölder Spaces. "International Journal of Development Research", Vol. 09, Issue. 05, pp. 27720-27730.
- Mamatov T (2019) Mapping Properties of Mixed Fractional Differentiation Operators in Hölder Spaces Defined by Usual Hölder Condition. Journal of Computer Science & Computational Mathematics,
- Kh. M. Murdaev, S.G Samko (1986) Weighted estimates of continuity modulus of fractional of functions having a prescribed continuity modulus with weight (Russian). Dep. in VINITI, Moscow, No. 3351-B, 42 p.
- Kh. M. Murdaev, S.G. Samko (1986) Mapping properties of fractioanl integro-differentiation in weighted  $\tilde{H}_0^\alpha(\rho)$  generalized Hölder spaces with the weight



- $\rho(x) = (x-a)^u (b-x)^v$  and given continuity modulus (Russian). Deponierte in VINITI, Moscow, No. 3350-B, 25 p.
15. B.S. Rubyn (1974) Fractional integrals in Hölder spaces with weight and operators of potential type. Izv. Akad. Nauk Armyan. SSR Ser. Mat. 9, No 4, pp. 308-324.
  16. B.S. Rubyn (1986) Fractional integrals and Riesz potentials with radial density in spaces with power weight. Izv. Akad. Nauk Armyan. SSR Ser. Mat. 21, No 5, pp. 488-503
  17. S.G. Samko, A.A. Kilbas and O.I. Marichev (1987) Fractional Integrals and Derivatives. Theory and Applications. Gordon and Breach. Sci. Publ., N. York - London, 1993, 1012 pp. (Russian Ed. - Fractional Intervals and Derivatives and Some of Their Applications, Nauka I Tekhnika, Minsk).
  18. S.G. Samko, H.M. Murdayev (1989) Weighted Zygmund estimates for fractional differentiation and integration and their applications. Trudy. Matem. Inst. Steklov 180 197-198; English transl. in: Proc. Steklov Inst. Math. (AMS), Issue 3 pp. 233-235.
  19. B. G. Vakulov (1986) Operators of potential type in generalized Hölder classes. Izv. universities. Mat. No. 11, pp.66-69.