

# Case Report: Unintentional Edible THC (Tetrahydrocannabinol) Ingestion (Gummi Worms) in a Child

James Espinosa MD<sup>1</sup>, Kelly Hamilton<sup>2</sup>, Alan Lucerna<sup>3</sup>, David Ho<sup>1</sup>, Henry Schuitema<sup>4</sup>

<sup>1</sup>Department of Emergency Medicine, Rowan University SOM/Jefferson - Stratford, NJ, USA

<sup>2</sup>Psychiatry Residency Program, Medstar-Georgetown University Hospital, Washington, DO

<sup>3</sup>Program Director, Emergency Medicine, Jefferson NJ/Rowan University SOM

<sup>4</sup>Chief, Department of Emergency Medicine, and Associate Chief Medical Officer, Jefferson, NJ

**\*Corresponding Author:** James Espinosa MD. Department of Emergency Medicine Rowan University SOM/Jefferson NJ 18 East Laurel Road

**Received date:** June 04, 2022; **Accepted date:** June 16, 2022; **Published date:** June 26, 2022

**Citation:** Espinosa J, Hamilton K, Lucerna A, Ho D, Schuitema H. (2022). Case Report: Unintentional Edible THC Ingestion (Gummi Worms) in a Child. International J. of Biomed Research. 2(8): DOI: 10.31579/IJBR-2022/084

**Copyright:** © 2022, Paul Andrew Bourne, this is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract:

We report the case of a 4-year-old male who presented to the Emergency Department (ED) with what his mother described as unusual fatigue and sleepiness. The mother related that family and friends had come to her house for a holiday gathering and that one of the guests had an accessible and open handbag that contained gummi worms with tetrahydrocannabinol (THC). The guest observed the child eating the gummi worms and immediately notified the patient's mother. Poison control was contacted from the ED. The child was observed for six hours and left the ED in good condition without any specific treatment needed. Awareness of the potential of unintentional edible cannabis in the pediatric population is important.

**Keywords:** Pediatric cannabinoid ingestion; pediatric THC ingestion

## Introduction

Pediatric inadvertent ingestions of THC infused candy is increasing. A review of acute cannabis toxicity notes that emergency department visits for cannabis toxicity in children increased from 1.2 per 100,000 population to 2.3 per 100,000 population. [1,2] This increase is seen both in young children and in adolescents. [3] Here we discuss the case of a six-year-old male with an inadvertent ingestion of THC-infused gummi worms

## The Case

We report the case of a 4-year male who presented to the ED with what his mother described as unusual fatigue and sleepiness. The mother related that family and friends had come to her house for a holiday gathering and that one of the guests had an accessible and open handbag that contained gummi worm THC. The guest observed the child eating the gummi worms and immediately notified the patient's mother. The child was initially alert and playful at the home but became increasingly fatigued and sleepy. The ingestion was one hour prior to ED presentation.

The mother brought the empty package of gummi worms to the ED. The bag was clearly marked that there were 10 gummi worms in a bag. Each

gummi contained 10 milligrams of THC. The child weighed 46 pounds (21 kilograms). The guest states that the bag had been opened and that one gummi had been ingested by the guest. Thus, the total maximum THC ingestion was 90 mg.

On arrival the child appeared tired and sleepy, but was easily awakened. Some mild tremor was noted on arrival. Vital signs on arrival were within normal limits for age. The oxygen saturation was 99%. There had been no nausea or vomiting. The patient was able to ambulate after the ingestion, but was in mother's arms on arrival.

Physical exam revealed a sleepy child who was easily awakened but then went back to sleep. He was able to follow commands easily. Physical exam was otherwise unremarkable. There were no focal findings. There was no nystagmus. There was no tremor.

Poison control was contacted. Poison control recommended 4 to 6 hours of observation in the ED and discharge if fully alert with normal vital signs and a normal exam. They did not recommend laboratory testing. A bedside glucose was within normal limits. The patient was reevaluated every 30 minutes. It was noted that the patient became fully alert after two hours in the ED. At three hours the patient was watching cartoons on a cell phone. The patient was ambulatory at 4 hours and wanted to play in

the hallways of the ED. The patient was discharged at 6 hours after arrival. Follow up contact with the patient's mom at 24 and 48 hours revealed that the patient showed full recovery and was playing, was alert and had a good appetite.

## Discussion

A review of acute cannabis toxicity notes that emergency department visits for cannabis toxicity in children increased from 1.2 per 100,000 population to 2.3 per 100,000 population. [1,2] This increase is seen both in young children and in adolescents. [3] Cannabis extracts can be infused into several common food items, such as brownies, cookies and candies (such as gummy worms) which can increase the risk of cannabis exposure to young children. [3] Cannabis has also been infused into popcorn and beverages. [4] Poison control calls for unintentional edible pediatric ingestions of cannabis have increased in the United States. [5]

Cannabis contains over 500 chemical components (cannabinoids) of which the most well described are tetrahydrocannabinol (THC) and cannabidiol (CBD). THC is the principal component responsible for central nervous system effects. [2]

THC can produce a wide range of symptoms. These are principally neurologic (euphoria, sedation, ataxia, stupor, coma); eye-related (nystagmus, injected conjunctiva, sluggish pupils); cardiovascular (tachycardia) and gastrointestinal (nausea, vomiting, increased appetite, thirst). [5] Cannabis can cause psychosis in children as well as hypothermia. [6]

Standard treatment is supportive. (airway, breathing and circulation) with treatment of specific symptomology. [2]

## Conclusion

ED visits for cannabis toxicity in children are increasing, both in young children and in adolescents. Cannabis extracts can be infused into several

common food items, such as brownies, cookies and candies (such as gummy worms) which can increase the risk of cannabis exposure to young children. THC can produce a wide range of symptoms. These are principally neurologic (euphoria, sedation, ataxia, stupor, coma); eye-related (nystagmus, injected conjunctiva, sluggish pupils); cardiovascular (tachycardia) and gastrointestinal (nausea, vomiting, increased appetite, thirst). Cannabis can cause psychosis in children as well as

hypothermia. Awareness of the potential of unintentional edible cannabis in the pediatric population is important.

## Conflicts of Interest

The authors declare that they have no conflicts of interest.

## References:

1. Levene RJ, Pollak-Christian E, Wolfram S. (2019). A 21st Century Problem: Cannabis Toxicity in a 13-Month-Old Child. *J Emerg Med.* 56(1):94-96.
2. Wong KU, Baum CR. (2019). Acute Cannabis Toxicity. *Pediatr Emerg Care.* 5(11):799-804.
3. Blohm E, Sell P, Neavyn M. (2019). Cannabinoid toxicity in pediatrics. *Curr Opin Pediatr.* 31(2):256-261.
4. Vo KT, Horng H, Li K, Ho RY, Wu AHB, Lynch KL, Smollin CG. (2018). Cannabis Intoxication Case Series: The Dangers of Edibles Containing Tetrahydrocannabinol. *Ann Emerg Med.* 2018 Mar;71(3):306-313.
5. Whitehill JM, Dilley JA, Brooks-Russell A, Terpak L, Graves JM. (2021). Edible Cannabis Exposures Among Children: 2017-2019. *Pediatrics.* 147(4).
6. Takakuwa KM, Schears RM. (2021). The emergency department care of the cannabis and synthetic cannabinoid patient: a narrative review. *Int J Emerg Med.* 10. 14(1):10.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

**Submit Manuscript**

DOI: [10.31579/IJBR-2022/084](https://doi.org/10.31579/IJBR-2022/084)

### Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://www.auctoresonline.org/journals/international-journal-of-biomed-research->