

Serotonergic modulation of visual processing

Amir-Hossein Darvish ^{1,2}, Ardeshir Nabizadeh ¹, Sahar Noursina ¹, Mir-Shahram Safari ^{1,2*}

¹Neuroscience Research Center, Shahid Beheshti University of Medical Sciences, Tehran 19615-1178, Iran.

²Brain Future Institute, Tehran, Iran.

***Corresponding Author:** Mir-Shahram Safari, Neuroscience Research Center, Shahid Beheshti University of Medical Sciences, Tehran 19615-1178, Iran.

Received date: August 30, 2024; **Accepted date:** septembre 13, 2024; **Published date:** septembre 20, 2024

Citation: M Amir-Hossein Darvish, Ardeshir Nabizadeh, Sahar Noursina, Mir-Shahram Safari, (2024), Serotonergic modulation of visual processing, *J. Neuroscience and Neurological Surgery*, 16(2); DOI:10.31579/2578-8868/336

Copyrights: ©, 2024, Mir-Shahram Safari. This is an open-access article distributed under the terms of The Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Abstract

Neuromodulators are chemicals that alter the activity of neural networks. They can have -immediate or delayed effects on neurons, affecting their properties at both nearby and distant synapses. Neuromodulators can influence neurons in various ways, allowing even small neural Networks generate a diverse range of functional outcomes. This diversity is crucial for the flexibility and adaptability of neural functions, including sensory processing.

Keywords: Neuromodulators ; Serotonergic modulation ; neural functions

Introduction

Neuromodulators are chemicals that alter the activity of neural networks. They can have -immediate or delayed effects on neurons, affecting their properties at both nearby and distant synapses. Neuromodulators can influence neurons in various ways, allowing even small neural Networks generate a diverse range of functional outcomes. This diversity is crucial for the flexibility and adaptability of neural functions, including sensory processing.

Serotonin is a widely conserved signaling molecule that influences various sensory systems across different species(1-3). It plays a crucial role in regulating states such as arousal, mood, and motivation (4-7). Serotonergic neurons, similar to other modulatory neurons, can influence a cell's activity without needing to form a direct synapse with it (8, 9). Sensory systems gather and process environmental information to create a internal representation of external world. Nevertheless, these systems must adapt to changes in the environment and the animal's internal state. This review aims to discuss how serotonin, a specific neuromodulator, influences sensory processing.

Serotonin as a modulator of sensory systems

Serotonin's effects on sensory systems have been extensively studied across different species and sensory modalities, making it an prime example for illustrating fundamental principles of neuromodulation. The sensory processing can modulated by serotonergic system in various behavioral scenarios. This influence can be complex and specific to particular stimuli(13). The diverse characteristics of serotonergic neurons allow serotonin to modulate sensory processing in a nonuniform and complex manner. This means that serotonin's effects can vary widely depending on the specific neurons and contexts involved.

Moreover, while the specific receptors involved are not fully understood, serotonin has been shown to reduce activity in proprioceptor and mechanosensory networks(14, 15). On the other hand, this neuromodulator also can increase excitability of photoreceptors by stimulation of 5-HT receptors (16-18), and sensitize these sensory afferents more than baseline. The influence of serotonin in V1, possibly via interactions with other neuromodulatory systems (19-23), might therefore aid visual processing during periods of quiet vigilance. This influence works by decreasing the spiking response gain to prevent an obvious orienting reaction in the animal, which is consistent with previously observed effects of serotonin in suppressing the acoustic startle response (24).

In early visual processing of cat, significantly reduced responses were observed after serotonin was applied iontophoretically in the lateral geniculate nucleus (LGN) (10) and V1 area of rat brain (11). In a more recent study involving awake macaques, where serotonin was iontophoretically applied and compared with the effects of pH-matched saline application, an overall reduction in sensory responses due to serotonin was observed (12).

Diversity of serotonergic neurons

DRN neurons have diverse pathways through which they send signals to other parts of the brain. Besides serotonin, these neurons can contain other neurotransmitters. The inherent electrical and chemical properties of these neurons vary. DRN neurons receive inputs from various sources, adding to their diversity. The genes expressed in these neurons differ, contributing to their functional diversity (13). Anatomical projections of Serotonergic system are heterogeneous, and this can reveal various

functional aspects within specific sensory systems. For example, serotonergic neurons do not distribute evenly across the insect visual system; instead, they selectively innervate different layers within each visual neuropil.(25-30).

Likewise, the extent of serotonergic innervation differs both intranuclear and internuclear in rodent auditory nuclei, such as those in the superior olive and the inferior colliculus. (31-33). Collectively, these facts demonstrate the diverse nature of serotonergic projections to networks that involved in sensory processing. This variability precise targeting of different processing layers or stimulus-specific subcircuits. Recent advances in technology have shown that subsets of serotonergic neurons are diverse in their molecular and anatomical characteristics(34-39). To investigate subsequent changes in V1, it is important to consider the extensive topographic organization within the DRN (40). Activation of 5-HT neurons may not only directly affect V1 but also modulate other cortical and subcortical regions that could, in turn, influence V1 activity (41). Moreover, it's important to note that, in addition to 5-HT neurons, some subpopulations of neurons that release glutamate in the raphe nuclei also express ePet. (42, 43).

Receptor Foundations of serotonergic Modulation

Serotonergic neurons themselves exhibit considerable diversity, and this is mirrored in the variety of serotonin receptors (5-HTRs). These receptors differ in their serotonin affinity, duration of action, and the secondary messenger systems they engage (44). The earliest 5-HTR appeared approximately over 500 million years ago (3), and vertebrates have seven primary 5-HTR families (5-HT1–7), while invertebrates possess at least three families (5-HT1, 2, and 7)(45).

Types of serotonin synapses and receptors involved in sensory modulation

Serotonergic projections to primary sensory areas are characterized by small, varicose axons that are broadly extended (46).The number of synaptic specializations, which are typically asymmetric, is minimal (47), indicating that serotonin primarily operates through bolus transmission from these varicosities. Although, there is ongoing debate regarding the reliance of neuromodulatory systems on "wired" transmission—highly localized and typically synaptic—or "volume" transmission, which is more spatially diffuse(48-51).

In the mammalian brain, seven serotonin receptor families, each with multiple subtypes, have been identified, contributing to the functional diversity of serotonin(52). While a comprehensive review of all receptors is beyond this scope, a few key receptors are worth noting. The 5-HT1A receptor is found on pyramidal neurons of cortex (53). In the primary visual cortex of macaques, the 5-HT1B and 5-HT2A receptors are most densely expressed, particularly in layer 4 (54). The 5-HT1B receptor is also highly expressed in the LGN but shows weak expression in other cortical regions, such as the auditory and somatosensory cortices (54).

In mice, GABAergic neurons that have the 5-HT3A receptor do not express the calcium-binding proteins parvalbumin or somatostatin, suggesting they may form a unique group of inhibitory interneurons (55). In anesthetized macaques, when examining the impact of the two most prominently expressed receptors in primary visual area, 5-HT1B and 5-HT2A, using receptor-specific ligands, researchers observed various pattern of reciprocal modulation for these receptors (54). In one study researchers found that when 5-HT was iontophoretically applied to the

visual cortical area of wakeful primates, it primarily reduced the gain of evoked responses at the population level while leaving ongoing activity unchanged.(12).

This distinct effect of 5-HT on gain response has been linked to the selective activation of 5-HT2A receptors (56) through several approaches, such as the subcutaneous injection of a hallucinogenic 5-HT2A receptor agonist in mice (57). The findings suggest that 5-HT induces separate inhibition of evoked and ongoing activity, influencing the gain of both in a divisive manner. simultaneous iontophoretic application of specific antagonists for 5-HT1A and 5-HT2A receptors (58-62) suggests that these receptors play distinct roles in regulating inhibition rate of ongoing and evoked network activity, respectively (63).

The findings demonstrates that activation of the serotonergic system influences two aspects of network activity in the visual cortex (V1): ongoing and evoked responses, with both being affected in a distinct and divisive manner. Each component is modulated through separate inhibitory effects of the 5-HT1A and 5-HT2A receptors, respectively. many studies in primary visual cortex, which used specific agonists for 5-HT1B and 5-HT2A receptors in combination with single-unit recordings in anesthetized monkeys, found bi-directional modulation depending on instantaneous firing rates. When a 5-HT2A agonist was applied, neurons with strong responses were suppressed, while those with weaker responses were facilitated, whereas the opposite occurred with the application of a 5-HT1B receptor agonist(54, 64).

Oppositely, the divisive scaling of stimulus driven responses is likely due to the dominant activation of excitatory 5-HT2A receptors, as inhibiting these receptors significantly lessened the inhibition of the evoked component. Furthermore, 5-HT2A receptors are linked to the Gq/11 signaling pathway (65), which typically increases neuronal firing rather than reducing it. This reduction in activity may be mediated by GABAergic neurons, which in turn decrease the activity of pyramidal neurons (66).

In fact, divisive modulation of responses in visual cortex has been shown to significantly depend on the activation of soma-targeting parvalbumin-expressing interneurons ((67), although other studies suggest differing roles (68, 69) for neurons that predominantly express 5-HT2A receptors (70, 71).

Activation of 5-HT2A receptors may also induce depolarizing currents in pyramidal neurons, leading to shunting inhibition, which increases conductance and affects both the gain and time constant of neuronal responses (72).

In two studies, selective activation of 5-HT2A receptors consistently produced a strong suppressive effect on the gain of visually evoked population responses (57, 73), despite cell-type and layer-specific differences across single cells (57). These findings support our current observations that DRN-triggered scaling of evoked responses is mediated by cortical 5-HT2A receptors at the population level. They also suggest that the distribution of a single neurotransmitter receptor type, or "receptome," can account for a distinct function in sensory processing (74-79). While 5-HT2A receptors regulate response gain, the scaling of ongoing V1 activity triggered by the DRN appears to be primarily controlled by 5-HT1A receptors (11).

Bidirectional interactions between serotonergic systems and sensory systems

Serotonergic neurons both influence sensory circuits and receive inputs from them. This bidirectional interaction allows the impact of serotonin on sensory processing to be modulated according to the sensory inputs the animal encounters.(45). This reciprocal interaction enables sensory networks to adjust to varying stimulus state. For example, some studies demonstrate that signal-to-noise ratio can be altered by serotonin (11, 80, 81), which helps maintain stable stimulus representation even in a noisy environment. Sensory input to serotonergic neurons can lead to specific modifications in serotonin release. This means that the release of serotonin can be tailored to the specific sensory stimuli being experienced. In certain instances, first sensory neurons and serotonergic cells are identical, enabling the sensory field to directly trigger releasing of serotonin (11, 80-87).

Sensory Stimulus-Driven Serotonin Release

Many serotonergic neurons, in addition to projecting to both sensory and nonsensory regions (36, 37, 88), also receive inputs from sensory systems. In certain cases, these neurons can acquire sensory input locally within the very networks they modulate. For example, in *Drosophila* and moths, CSDn activity is affected by odors (89-92) through direct synaptic connections with antennal lobe principal neurons (91, 93-95). The pattern of local input to a single neuron can differ across sensory networks, as CSDns can be both stimulated and suppressed by the same odor due to local synaptic inputs targeting various neuronal compartments. (90).

Serotonergic neurons in the DRN and MRN also receive inputs from various cortical and subcortical sensory areas, such as the inferior and superior colliculi and brainstem sensory nuclei (96, 97). Moreover, these neurons are responsive to stimuli across various sensory modalities (98-102). Although the strongest sensory responses are often found in nonserotonergic raphe neurons, serotonergic neurons themselves have been shown to be responsive to sensory inputs (103-105), and nonserotonergic DRN neurons may relay indirect sensory input to serotonergic neurons. Furthermore, some sensory responses in serotonergic DRN neurons exhibit very short latencies, suggesting input from early stages of sensory processing (103). Consequently, whether they are confined to a single network or span multiple networks, serotonergic neurons, by having close access to the history of network activity, can adjust their modulatory effects based on the current stimulus or circuit state (106).

Serotonergic Neurons Adjust Based on Behavioral State and Context

The circumstances surrounding serotonin release are multifaceted, encompassing both dependent and independent stimulus state. Researches shows that serotonergic neurons react to sensory stimuli but are also affected by factors like internal states, movement, and the importance of sensory events in relation to past experiences. This implies that serotonin helps transmit information to sensory systems regarding both the external environment and the internal conditions in which sensory events happen (45).

By using varying stimulus intensities, the normalization of visual responses was identified. In anesthetized animals, the ongoing activity contributed to response normalization by acting as a subtractive factor. However, in the awake state, normalization was governed by the gain of the evoked response and was independent of any concurrent suppression of ongoing activity.

Inhibition tends to dominate activity in the awake cortex (107), possibly alongside elevated 5-HT levels (108, 109). Consequently, since the inhibition of baseline activity depends on initial baseline levels, the additional impact of the spontaneous part on normalization in the anesthetized condition could be due to significant fluctuations in its amplitude, which are less common during wakefulness (107). As a result, the extent of serotonin-regulated integration of sensory input and Baseline activity dependent to cortical condition and baseline levels of serotonin, and this seems to be predominantly controlled by 5-HT1A receptors (110).

In summary, the interplay between 5-HT1A and 5-HT2A receptors leads to a notable and significant adjustment in both spontaneous and evoked components of population activity in V1. One key observation is that the inhibition of gain modulation induced by 5-HT is more pronounced in the anesthetized condition compared to the awake state. This suggests that, while awake, response normalization depends less on spontaneous activity and is less affected by internal cortical signals.

Given that spontaneous activity reflects top-down expectations, whereas Neural responses to stimuli provide bottom-up sensory information, any imbalance in the activation of these receptors—whether due to specific agonist application or irregular receptor expression—can disrupt the integration of these components and thus affect cortical information processing. Such an imbalance could lead to an excessive focus on internally generated expectations(111, 112) at the expense of sensory input, or the reverse (113).

Serotonergic Modulation of Sensory Computations

Sensory systems use various strategies to dynamically adjust the range of individual stimulus features they encode. Given the widespread presence of serotonergic systems, it's unsurprising that serotonin either modulates or directly participates in these processes. For example, as animals move through their environment, they encounter fluctuations in stimulus intensity. In situations where sensory input is intense (such as bright light or strong odors), neurons may struggle to properly encode the stimulus due to saturation. On the other hand, when stimuli are at low intensities, such as the faint scent of a predator, animals may fail to detect them. To address these challenges, sensory systems typically employ a range of computations, including "gain control" (72), which adaptively adjusts the input-to-output ratio of a network. Across different sensory modalities, 5-HT receptors (5-HTRs) expressed by sensory afferents enable direct serotonergic modulation of sensory input gain(45).

Serotonin can decrease the signal-to-noise ratio by reducing evoked activity more than spontaneous activity (11) The concept of serotonergic modulation of population codes is relatively recent, and further research in this area could help bridge the gap between the effects of monoamines on neural codes and sensory-related behaviors (114). This modulation involves a reduction in retinotectal transmission mediated by 5HT1B receptors selectively expressed by optic afferents, though postsynaptic 5HT1A receptors may also contribute to a decline in responsiveness (115).

In macaque V1, gain changes are the most frequently reported modulatory effect. Both acetylcholine (ACh) and serotonin (and likely norepinephrine, NA) modify gain in some way; however, studies vary (and possibly the systems themselves) in terms of how many neurons are suppressed versus enhanced. Other reviews have presented studies on ACh and serotonin (12, 54), that report conflicting findings regarding the

dominance of enhancement versus suppression across neurons. Notably, serotonin does not alter rate variability or noise correlations in macaque V1 (12).

The reported effects of serotonin on sensory processing have been inconsistent (11, 54, 113, 114, 116) making it difficult to develop a straightforward computational explanation. However, studies showed that across the neuronal population in V1 and across different stimulus dimensions, serotonergic modulation is surprisingly simple: serotonin primarily reduces the gain of visual responses, with minimal impact on tuning properties. Results suggest that serotonin is well-suited to controlling the response gain of neurons in V1, potentially complementing existing gain control mechanisms.

Across various visual stimulus dimensions, serotonergic modulation was uniformly characterized by a reduction in response gain, a slight slowing of response dynamics, and no systematic changes in neuronal variability, co-variability, or stimulus selectivity.

These effects could be captured by a model in which serotonin induces a simple additive change in the threshold-linear spiking nonlinearity. Overall, the observed modulation was homogeneous, resulting in a straightforward decrease in response gain across the neural population (12). Such gain modulation is an essential aspect of cortical computation (117) allowing responses to be modulated without altering receptive field properties, and making it well-suited to adjust responses according to the animal's internal state, influenced by the context's valence (118).

The inhibitory effect of serotonin was the predominant pattern observed across the large neuronal population. This reduction was mainly attributed to a multiplicative change (gain change) in the neuronal tuning curves (119). Behaviorally, a recent study that administered a serotonin-reuptake inhibitor to enhance serotonin's effect during a color discrimination task in macaques found that reaction times slowed and perceptual performance deteriorated (120), consistent with the expected outcome of reduced visual responses.

On the other hand, a reduction in the gain of spontaneous responses (113), aligns with an increased signal-to-noise ratio (SNR), or a reduction in the gain of tuning curves (116).

The mechanisms and nature of how specific populations of neuromodulatory inputs influence sensory processing content remain largely unclear.

Studies shown that Serotonin can modulate the transmission of early visual information within critical regions, such as the dorsolateral geniculate nucleus (dLGN) (121, 122), the suprachiasmatic nucleus (123, 124), and the superior colliculus (115), in addition to its effects on early olfactory and auditory information (116, 125).

Serotonergic modulation of retinal processing

Specifically, in vitro pharmacological administering high dosage of serotonin or its agonists has been shown to reduce retinal axon stimulation-evoked glutamate release via presynaptically expressed 5-HT1B receptors (121-123). However, it remains uncertain whether "endogenous" serotonin release significantly modulates the activity of mouse retinal axonal boutons in vivo, and if this modulation is selective for particular retinal axons.

An Study indicates that serotonin preferentially suppresses retinal ganglion cell (RGC) axonal boutons with high baseline activity that

respond to full-field stimuli. This selective suppression by serotonin may alter the tuning of the postsynaptic cell toward smaller, more localized stimuli, similar to what has been observed in zebrafish (126).

In fact, electrical stimulation of the dorsal raphe nucleus (DRN) selectively diminishes evoked activity in thalamocortical neurons with large receptive fields (127), which aligns with observations of stronger suppression of full-field boutons. Also slow fluctuations in serotonin release within the dLGN, a small fraction of which could be attributed to a weak anti-correlation with arousal levels (128). The behavioral contexts that drive these serotonin fluctuations are still largely unknown, as are the roles of serotonergic suppression of retinothalamic transmission in shaping downstream visual processing and behavior (128). However, while serotonin preferentially suppresses RGC axons that are strongly responsive to full-field luminance changes, arousal more selectively suppresses boutons that are more responsive to localized stimuli. This suggests the intriguing possibility that serotonergic axons and other modulatory inputs may implement multiple, complementary, state-dependent selective filters for specific visual information channels at a critical bottleneck in the pathway, before these channels reach thalamocortical neurons and are further relayed and amplified in brain regions responsible for guiding behavior and learning (128).

Serotonergic modulation of visual processing by various receptors

The 5-HT1B receptor has been shown to influence neurotransmission in various pathways, including the retinocollicular (129) retino-suprachiasmatic nuclear (123), and thalamocortical (130) pathways. Given the increased expression of 5-HT1B and 5-HT2A receptor genes in V1, it is plausible that these two receptors play significant roles in modulating neurotransmission in this region (54).

Activity-dependent expression of 5-HT1B receptor mRNA has also been observed in the LGN. To determine the time needed for monocular inactivation to take effect, shorter durations of monocular inactivation (1 day, 6 hours, and 3 hours) were examined. The downregulation of 5-HT1B and 5-HT2A receptor mRNAs in V1 was detected even after just 3 hours of monocular inactivation. These findings suggest that the specific expression patterns of 5-HT1B and 5-HT2A receptor mRNAs in V1 are maintained by ongoing visual activity. Consequently, it is speculated that 5-HT1B agonist (CP93129) mainly facilitated visual responses in V1 neurons, although it tended to suppress neurons with low firing rates. This example illustrates that activation of 5-HT1B receptors enhances responses to high-contrast (>30%) stimulation but is either suppressive or ineffective for low-contrast stimuli. The results indicate that the effect of the 5-HT1B receptor agonist is contingent on each neuron's response level and that it improves the signal-to-noise ratio of visual input from the LGN to the cortex (54).

In vivo electrophysiological experiments demonstrated that 5-HT1B and 5-HT2A agonists modulate V1 neuron responses in macaque monkeys. This analysis revealed that 5-HT2 agonist (DOI) facilitates visual responses in neurons with low firing rates but suppresses those with high firing rates (54).

They conclude that 5-HT2A receptors exhibit response-dependent modulatory effects, but their effects are opposite to those of 5-HT1B receptors. Then, they indicate that the laminar distributions of the effects of 5-HT1B agonist (CP93129), DOI, and 5-HT2A antagonist (ketanserin). This contrasts with the highly layer-specific distribution of 5-HT1B and 5-HT2A receptor mRNAs (54).

Seeburg et al.(122) shown that each receptor can exert both suppressive and facilitative effects depending on the firing rate of the recorded neurons. Regarding the 5-HT1B receptor, similar context-dependent bidirectional modulation has been observed in vitro using brain slices containing the optic tract and LGN (122) or the ventral posterior medial nucleus of the thalamus and somatosensory cortex (130). For instance, Seeburg et al. in 2004 (122) found that 5-HT1B receptor-mediated serotonergic modulation of LGN neuron responses to optic tract stimulation depends on the stimulus's temporal frequency.

The 5-HT1 receptor agonist suppresses retinogeniculate transmission for low-frequency inputs but is either ineffective or facilitative for high-frequency inputs. The authors suggested that alleviation of synaptic depression due to high-frequency stimulation might underlie these effects of the 5-HT1B receptors.

On the other hand, activation of 5-HT2A receptors is known to exert direct facilitatory actions on pyramidal neurons and interneurons (131, 132), indirectly inhibiting neighboring pyramidal neurons (133-135). Thus, serotonin likely has complex effects by regulating the relative activity of excitatory and inhibitory neurons within local circuits through 5-HT1B and 5-HT2A receptors (54).

Serotonin release in V1 could be locally regulated by 5-HT1B autoreceptors on the presynaptic terminals of raphe neurons that project to V1 or potentially by prefrontal cortex activity (136). Therefore, serotonin's cortical effects likely depend on the dynamic regulation of its levels and receptors. Although the enhanced expression of 5-HT1B receptor mRNA in the LGN and V1 suggests its primary role in the visual system, it is also widely expressed in the thalamus. In a study, it was demonstrated that activating 5-HT1B receptors in V1 generally enhances visual responses but tends to suppress weak responses. This suggests that, in geniculocortical transmission, non-synchronized spontaneous activity (noise) from LGN neurons is reduced by 5-HT1B receptor-mediated suppression, while visually evoked synchronized signals are preserved or efficiently transmitted to V1, thereby enhancing the signal-to-noise ratio in the input-output relationship.

Conversely, neurons in V1's input layers, which express the 5-HT2A receptor abundantly, may act as gain controllers by enhancing weak signal responses and suppressing excessive responses. Therefore, we suggest that serotonin release in V1 exerts coordinated modulatory effects through 5-HT1B and 5-HT2A receptors on V1 neurons. It is thus possible that the serotonin system has contributed to the evolution of the primate visual system's sophisticated function (54).

In a study, it was investigated visual processing and experience-dependent learning in SERT-deficient mice. They did not find significant alterations in orientation, spatial frequency, and contrast tuning in naive mice, consistent with a previous operant conditioning study that found intact learning in visual discrimination tasks in SERT-deficient mice (137). Additionally, compensatory mechanisms may partially correct for the lack of functional SERT to maintain cortical development (138). However, they observed a lack of bias toward cardinal orientations in V1 of SERT-deficient mice before visual experience, which partially recovered in SERT heterozygous mice after perceptual experience but not in knockout animals. Observations of prolonged oscillatory activity following perceptual experience in SERT-deficient mice further support the hypothesis that 5-HT may play a role in cortical plasticity. They also found decreased orientation selectivity and broadened orientation tuning

in SERT knockout mice. Interestingly, these changes resemble those seen in Fmr1 knockout mice, which are known to result from hypoactivation of parvalbumin-positive fast-spiking interneurons and corresponding circuit alterations (139, 140).

Heterogenous actions of serotonin on interneurons in visual cortex

It is interesting that serotonergic modulation of inhibitory neurons is cell-type specific. Xiang and Prince (141) suggest that serotonergic activation exerts complex influences on cortical inhibitory networks, potentially leading to alterations in cortical information processing (141). The impact of serotonin (5-HT) on the excitability of two cortical interneuron subtypes, fast-spiking (FS) and low-threshold spiking (LTS) cells, as well as on spontaneous inhibitory postsynaptic currents (sIPSCs) in layer V pyramidal cells, was examined in rat visual cortical slices using whole-cell recording methods (141). It was observed that the application of 5-HT induced excitation in half of the FS cells and a small portion of LTS cells, while it caused inhibition in roughly half of the FS cells and the majority of LTS cells. In a few FS and LTS cells, serotonin application triggered excitation followed by inhibition (141).

Serotonergic modulation of spatial attention and receptive field

Patel et al. (142) explore the effects of serotonergic modulation on local network processing in the primary visual cortex (V1) of awake macaques. Their research provides valuable insights into how serotonin influences sensory processing, drawing parallels with the effects of spatial attention. The authors highlight that serotonin application leads to a reduction in local field potential (LFP) power and spike-field coherence, indicating decreased functional connectivity within the local network.

This reduction in synchronization is thought to enhance the signal-to-noise ratio, thereby improving the efficiency of sensory processing. Interestingly, these effects mirror those observed with spatial attention, suggesting that serotonin may play a similar role in modulating sensory inputs.

Patel et al.(142) propose that serotonin's modulation of V1 contributes to a state of "quiet vigilance," where the brain remains alert but less reactive to distracting stimuli. This state is beneficial for maintaining focus on relevant visual inputs while minimizing the impact of irrelevant stimuli that this mechanism allows for a more stable and focused perception, which is crucial for tasks requiring sustained attention. The study also discusses the broader implications of serotonergic modulation across different sensory modalities.

The authors speculate that the effects observed in V1 might extend to other sensory systems, highlighting serotonin's role in fine-tuning sensory processing and maintaining perceptual stability. This cross-modal influence underscores the importance of serotonin in optimizing sensory perception and attention. Their study provides valuable evidence that serotonin, much like spatial attention, can shape sensory experiences by modulating local network dynamics. This insight is crucial for our understanding of how neuromodulators influence sensory modalities, particularly vision.

Serotonin (5-HT) generally diminished the effectiveness of synaptically mediated excitation and inhibition, resulting in a reduction or complete loss of neuronal responsiveness to visual stimuli. Regarding the effects of 5-HT on receptive field characteristics, a simple difference in threshold versus subthreshold synaptic inputs appears insufficient to explain cases where the area of the visual field that could activate a cell significantly

changed during 5-HT application. The reduction in evoked response magnitude observed in many instances could be attributed to the suppressive influence of 5-HT. However, the observed shift in the receptive field within visual space, likely involves a more intricate reconfiguration of synaptic balances that establish receptive field boundaries (11). Serotonin (5-HT) inhibits stimulus-evoked neuronal firing in the sensory regions of the cerebral cortex (11, 143, 144), while having a lesser impact on the background activity of these neurons. Additionally, 5-HT reversibly modifies the receptive fields of visual cortical neurons, causing a shift in the visual field towards the temporal zone(11).

Future directions:

Serotonergic system seems to be one of complicated brain modulatory systems. Different kind of serotonin receptors, difference in serotonergic neurons properties and non-homogenous projection to different sensory areas, are some of the complex aspects of this system. Considering role of before mentioned factors in various time points of visual processing in a cell-type and layer/area specific manner in multiple aspects of vision are necessary. These studies would be possible by more advanced neuroscience techniques with micro/nano meter spatial resolution and micro/nano second temporal resolution of recording of activities of many neurons, in multiple areas of animal brain, in vivo.

References

1. Gaspar P, Lillesaar CJPTotRSBBS. (2012). Probing the diversity of serotonin neurons. 367(1601):2382-2394.
2. Hay-Schmidt AJPotRSolSBBS. (2000). The evolution of the serotonergic nervous system. 267(1448):1071-1079.
3. Peroutka S, Howell TJN. (1994). The molecular evolution of G protein-coupled receptors: focus on 5-hydroxytryptamine receptors. 33(3-4):319-324.
4. Cools R, Roberts AC, Robbins TWJTics. (2008). Serotonergic regulation of emotional and behavioural control processes. 12(1):31-40.
5. Luo M, Li Y, Zhong WJNol, memory. (2016). Do dorsal raphe 5-HT neurons encode "beneficialness"? 135:40-49.
6. Miyazaki KW, Miyazaki K, Doya KJEJoN. (2011). Activation of the central serotonergic system in response to delayed but not omitted rewards. 33(1):153-160.
7. Monti JMJSmr. (2011). Serotonin control of sleep-wake behavior. 15(4):269-281.
8. Eid L, Champigny MF, Parent A, Parent MJEJoN. (2013). Quantitative and ultrastructural study of serotonin innervation of the globus pallidus in squirrel monkeys. 37(10):1659-1668.
9. Fuxe K, Agnati LF, Marcoli M, Borroto-Escuela DOJNr. (2015). Volume transmission in central dopamine and noradrenaline neurons and its astroglial targets. 40:2600-2614.
10. Phillis J, Teb  cis AJTJoP. (1967). The responses of thalamic neurones to iontophoretically applied monoamines. 192(3):715-745.
11. Waterhouse BD, Azizi SA, Burne RA, Woodward DJJBr. (1990). Modulation of rat cortical area 17 neuronal responses to moving visual stimuli during norepinephrine and serotonin microiontophoresis. 514(2):276-292.
12. Seillier L, Lorenz C, Kawaguchi K, Ott T, Nieder A, Pourriahi P, et al. (2017). Serotonin decreases the gain of visual responses in awake macaque V1. 37(47):11390-11405.
13. Okaty BW, Commons KG, Dymecki SMJNrn. (2019). Embracing diversity in the 5-HT neuronal system. 20(7):397-424.
14. Gaudry Q, Kristan Jr WBJNn. (2009). Behavioral choice by presynaptic inhibition of tactile sensory terminals. 12(11):1450-1457.
15. Nagata A, Nakayama K, Nakamura S, Mochizuki A, Gemba C, Aoki R, et al. (2019). Serotonin1B receptor-mediated presynaptic inhibition of proprioceptive sensory inputs to jaw-closing motoneurons. 149:260-267.
16. Cheng KY, Frye MAJJoCPA. (2020). Neuromodulation of insect motion vision. 206(2):125-137.
17. Han L, Zhong Y-M, Yang X-LJBrb. (2007). 5-HT2A receptors are differentially expressed in bullfrog and rat retinas: a comparative study. 73(4-6):273-277.
18. Pootanakit K, Prior KJ, Hunter DD, Brunken WJJVn. (1999). 5-HT2a receptors in the rabbit retina: potential presynaptic modulators. 16(2):221-230.
19. Dayan PJN. (2012). Twenty-five lessons from computational neuromodulation. 76(1):240-256.
20. Flavell SW, Gogolla N, Lovett-Barron M, Zelikowsky MJN. (2022). The emergence and influence of internal states. 110(16):2545-2570.
21. Krueger J, Disney AAJJoCN. (2019). Structure and function of dual-source cholinergic modulation in early vision. 527(3):738-750.
22. L  rincz ML, Adamantidis ARJPIn. (2017). Monoaminergic control of brain states and sensory processing: existing knowledge and recent insights obtained with optogenetics. 151:237-253.
23. Marder EJN. (2012). Neuromodulation of neuronal circuits: back to the future. 76(1):1-11.
24. Davis M, Strachan DI, Kass EJS. (1980). Excitatory and inhibitory effects of serotonin on sensorimotor reactivity measured with acoustic startle. 209(4455):521-523.
25. Hamanaka Y, Kinoshita M, Homberg U, Arikawa KJPo. (2012). Immunocytochemical localization of amines and GABA in the optic lobe of the butterfly, *Papilio xuthus*. 7(7):e41109.
26. Homberg U, Hildebrand JJJoCN. (1989). Serotonin immunoreactivity in the optic lobes of the sphinx moth *Manduca sexta* and colocalization with FMRFamide and SCPB immunoreactivity. 288(2):243-253.
27. N  ssel D, Ohlsson L, Sivasubramanian PJTJocn. (1987). Postembryonic differentiation of serotonin-immunoreactive neurons in fleshfly optic lobes developing in situ or cultured in vivo without eye discs. 255(3):327-340.
28. Paulk AC, Phillips-Portillo J, Dacks AM, Fellous J-M, Gronenberg WJJJoN. (2008). The processing of color, motion, and stimulus timing are anatomically segregated in the bumblebee brain. 28(25):6319-6332.
29. Sch  rmann F, Klemm NJJoCN. (1984). Serotonin-immunoreactive neurons in the brain of the honeybee. 225(4):570-580.
30. Vall  s AM, White KJJJoCN. (1988). Serotonin-containing neurons in *Drosophila melanogaster*: Development and distribution. 268(3):414-428.

31. Hurley LM, Thompson AMJJoCN. (2001). Serotonergic innervation of the auditory brainstem of the Mexican free-tailed bat, *Tadarida brasiliensis*. 435(1):78-88.
32. Keesom SM, Morningstar MD, Sandlain R, Wise BM, Hurley LMJBr. (2018). Social isolation reduces serotonergic fiber density in the inferior colliculus of female, but not male, mice. 1694:94-103.
33. Thompson GC, Thompson AM, Garrett KM, Britton BHJOH, Surgery N. (1994). Serotonin and serotonin receptors in the central auditory system. 110(1):93-102.
34. Calizo LH, Akanwa A, Ma X, Pan Y-z, Lemos JC, Craige C, et al. (2011). Raphe serotonin neurons are not homogenous: electrophysiological, morphological and neurochemical evidence. 61(3):524-543.
35. Fernandez SP, Cauli B, Cabezas C, Muzerelle A, Poncer J-C, Gaspar PJBS, et al. (2016). Multiscale single-cell analysis reveals unique phenotypes of raphe 5-HT neurons projecting to the forebrain. 221:4007-4025.
36. Huang KW, Ochandarena NE, Philson AC, Hyun M, Birnbaum JE, Cicconet M, et al. (2019). Molecular and anatomical organization of the dorsal raphe nucleus. 8:e46464.
37. Ren J, Isakova A, Friedmann D, Zeng J, Grutzner SM, Pun A, et al. (2019). Single-cell transcriptomes and whole-brain projections of serotonin neurons in the mouse dorsal and median raphe nuclei. 8:e49424.
38. Spaethling JM, Piel D, Dueck H, Buckley PT, Morris JF, Fisher SA, et al. (2014). Serotonergic neuron regulation informed by in vivo single-cell transcriptomics. 28(2):771.
39. Templin JS, Bang SJ, Soiza-Reilly M, Berde CB, Commons KGJBr. (2012). Patterned expression of ion channel genes in mouse dorsal raphe nucleus determined with the Allen Mouse Brain Atlas. 1457:1-12.
40. Muzerelle A, Scotto-Lomassese S, Bernard JF, Soiza-Reilly M, Gaspar PJBS, Function. (2016). Conditional anterograde tracing reveals distinct targeting of individual serotonin cell groups (B5–B9) to the forebrain and brainstem. 221:535-561.
41. Gilbert CD, Li WJNrn. (2013). Top-down influences on visual processing. 14(5):350-363.
42. McDevitt RA, Tiran-Cappello A, Shen H, Balderas I, Britt JP, Marino RA, et al. (2014). Serotonergic versus nonserotonergic dorsal raphe projection neurons: differential participation in reward circuitry. 8(6):1857-1869.
43. Sos KE, Mayer MI, Cserép C, Takács FS, Szőnyi A, Freund TF, et al. (2017). Cellular architecture and transmitter phenotypes of neurons of the mouse median raphe region. 222:287-299.
44. Nichols DE, Nichols CDJCr. (2008). Serotonin receptors. 108(5):1614-1641.
45. Sizemore TR, Hurley LM, Dacks AMJJon. (2020). Serotonergic modulation across sensory modalities. 123(6):2406-2425.
46. Hornung J-P. (2010). The neuronatomy of the serotonergic system. Handbook of Behavioral Neuroscience. 21: Elsevier; p. 51-64.
47. Descarries L, Riad M, Parent M. (2010). Ultrastructure of the serotonin innervation in the mammalian central nervous system. Handbook of behavioral neuroscience. 21: Elsevier; p. 65-101.
48. Rice ME, Cragg SJJBr. (2008). Dopamine spillover after quantal release: rethinking dopamine transmission in the nigrostriatal pathway. 58(2):303-313.
49. Sarter M, Parikh V, Howe WMJNRN. (2009) Phasic acetylcholine release and the volume transmission hypothesis: time to move on. 10(5):383-390.
50. Fuxe K, Borroto-Escuela DOJNrr. (2016). Volume transmission and receptor-receptor interactions in heteroreceptor complexes: understanding the role of new concepts for brain communication. 11(8):1220-1223.
51. Sulzer D, Cragg SJ, Rice MEJBG. (2016). Striatal dopamine neurotransmission: regulation of release and uptake. 6(3):123-148.
52. Mengod G, Cortés R, Vilaró MT, Hoyer D. (2010). Distribution of 5-HT receptors in the central nervous system. Handbook of behavioral neuroscience. 21: Elsevier; p. 123-138.
53. DeFelipe J, Arellano JI, Gómez A, Azmitia EC, Muñoz AJJoCN. (2001) Pyramidal cell axons show a local specialization for GABA and 5-HT inputs in monkey and human cerebral cortex. 433(1):148-155.
54. Watakabe A, Komatsu Y, Sadakane O, Shimegi S, Takahata T, Higo N, et al. (2009). Enriched expression of serotonin 1B and 2A receptor genes in macaque visual cortex and their bidirectional modulatory effects on neuronal responses. 19(8):1915-1928.
55. Rudy B, Fishell G, Lee S, Hjerling-Leffler JJDn. (2011). Three groups of interneurons account for nearly 100% of neocortical GABAergic neurons. 71(1):45-61.
56. Zhang G, Stackman Jr RWJFip. (2015). The role of serotonin 5-HT2A receptors in memory and cognition. 6:225.
57. Michaiel AM, Parker PR, Niell CMJCr. (2019). A hallucinogenic serotonin-2A receptor agonist reduces visual response gain and alters temporal dynamics in mouse V1. 26(13):3475-3483. e4.
58. Dyck RH, Cynader MSJPotNAoS. (1993). An interdigitated columnar mosaic of cytochrome oxidase, zinc, and neurotransmitter-related molecules in cat and monkey visual cortex. 90(19):9066-9069.
59. Jakab RL, Goldman-Rakic PSJPotNAoS. (1998). 5-Hydroxytryptamine2A serotonin receptors in the primate cerebral cortex: possible site of action of hallucinogenic and antipsychotic drugs in pyramidal cell apical dendrites. 95(2):735-740.
60. Leysen JJCDT-C, Disorders N. (2004). 5-HT2 receptors. 3(1):11-26.
61. Riga MS, Bortolozzi A, Campa L, Artigas F, Celada PJN. (2016). The serotonergic hallucinogen 5-methoxy-N, N-dimethyltryptamine disrupts cortical activity in a regionally-selective manner via 5-HT1A and 5-HT2A receptors. 101:370-378.
62. Shukla R, Watakabe A, Yamamori TJFinc. (2014). mRNA expression profile of serotonin receptor subtypes and distribution of serotonergic terminations in marmoset brain. 8:52.
63. Azimi Z, Spoida K, Barzan R, Wollenweber P, Mark MD, Herlitze S, et al. (2018). Subtraction and division of visual cortical population responses by the serotonergic system. 444943.

64. Shimegi S, Kimura A, Sato A, Aoyama C, Mizuyama R, Tsunoda K, et al. (2016). Cholinergic and serotonergic modulation of visual information processing in monkey V1. 110(1-2):44-51.
65. Hannon J, Hoyer DJBr. (2008). Molecular biology of 5-HT receptors. 195(1):198-213.
66. Gellman RL, Aghajanian GKJBr. (1993). Pyramidal cells in piriform cortex receive a convergence of inputs from monoamine activated GABAergic interneurons. 600(1):63-73.
67. Wilson NR, Runyan CA, Wang FL, Sur MJN. (2012). Division and subtraction by distinct cortical inhibitory networks in vivo. 488(7411):343-348.
68. Lee S-H, Kwan AC, Zhang S, Phoumthipphavong V, Flannery JG, Masmanidis SC, et al. (2012). Activation of specific interneurons improves V1 feature selectivity and visual perception. 488(7411):379-383.
69. Seybold BA, Phillips EA, Schreiner CE, Hasenstaub ARJN. (2015). Inhibitory actions unified by network integration. 87(6):1181-1192.
70. Puig MV, Gullledge ATJMN. (2011). Serotonin and prefrontal cortex function: neurons, networks, and circuits. 44:449-464.
71. Weber ET, Andrade RJFin. (2010). Htr2a gene and 5-HT2A receptor expression in the cerebral cortex studied using genetically modified mice. 4:36.
72. Carandini M, Heeger DJJNrn. (2012). Normalization as a canonical neural computation. 13(1):51-62.
73. Eickelbeck D, Karapinar R, Jack A, Suess ST, Barzan R, Azimi Z, et al. (2019). CaMello-XR enables visualization and optogenetic control of Gq/11 signals and receptor trafficking in GPCR-specific domains. 2(1):60.
74. Akimova E, Lanzenberger R, Kasper SJBp. (2009). The serotonin-1A receptor in anxiety disorders. 66(7):627-635.
75. Carhart-Harris RL, Muthukumaraswamy S, Roseman L, Kaalen M, Droog W, Murphy K, et al. (2016). Neural correlates of the LSD experience revealed by multimodal neuroimaging. 113(17):4853-4858.
76. Deco G, Cruzat J, Cabral J, Knudsen GM, Carhart-Harris RL, Whybrow PC, et al. (2018). Whole-brain multimodal neuroimaging model using serotonin receptor maps explains non-linear functional effects of LSD. 28(19):3065-3074. e6.
77. Goldberg HL, Finnerty RJTAjop. (1979). The comparative efficacy of buspirone and diazepam in the treatment of anxiety. 136(9):1184-1187.
78. González-Maeso J, Weisstaub NV, Zhou M, Chan P, Ivic L, Ang R, et al. (2007). Hallucinogens recruit specific cortical 5-HT2A receptor-mediated signaling pathways to affect behavior. 53(3):439-452.
79. Tauscher J, Bagby RM, Javanmard M, Christensen BK, Kasper S, Kapur SJAJoP. (2001). Inverse relationship between serotonin 5-HT1A receptor binding and anxiety: a [11C] WAY-100635 PET investigation in healthy volunteers. 158(8):1326-1328.
80. Hurley LM, Pollak GDJJoN. (1999). Serotonin differentially modulates responses to tones and frequency-modulated sweeps in the inferior colliculus. 19(18):8071-8082.
81. Waterhouse BD, Moises HC, Woodward DJJBrb. (1986). Interaction of serotonin with somatosensory cortical neuronal responses to afferent synaptic inputs and putative neurotransmitters. 17(4):507-518.
82. Iwanir S, Brown AS, Nagy S, Najjar D, Kazakov A, Lee KS, et al. (2016). Serotonin promotes exploitation in complex environments by accelerating decision-making. 14:1-15.
83. Liu H, Qin L-W, Li R, Zhang C, Al-Sheikh U, Wu Z-XJPotNAoS. (2019). Reciprocal modulation of 5-HT and octopamine regulates pumping via feedforward and feedback circuits in *C. elegans*. 116(14):7107-7012.
84. Shao J, Zhang X, Cheng H, Yue X, Zou W, Kang LJPA-EJoP. (2019). Serotonergic neuron ADF modulates avoidance behaviors by inhibiting sensory neurons in *C. elegans*. 471:357-363.
85. Watanabe H, Shimohigashi M, Yokohari FJJoCN. (2014). Serotonin-immunoreactive sensory neurons in the antenna of the cockroach *Periplaneta americana*. 522(2):414-434.
86. Nielsen K, Brask D, Knudsen GM, Aznar SJS. (2006). Immunodetection of the serotonin transporter protein is a more valid marker for serotonergic fibers than serotonin. 59(5):270-276.
87. Vyas P, Wu JS, Jimenez A, Glowatzki E, Fuchs PAJSr. (2019). Characterization of transgenic mouse lines for labeling type I and type II afferent neurons in the cochlea. 9(1):5549.
88. Gaudry QJTYJoB, Medicine. (2018). Focus: Sensory Biology and Pain: Serotonergic Modulation of Olfaction in Rodents and Insects. 91(1):23.
89. Hill ES, Iwano M, Gatellier L, Kanzaki R. (2002). Morphology and Physiology of the Serotonin-immunoreactive Putative Antennal Lobe Feedback Neuron in the Male Silkworm *Bombyx mori*. Chemical Senses. 27(5):475-483.
90. Zhang X, Coates K, Dacks A, Gunay C, Scott Lauritzen J, Li F, et al. (2019). Regulation of modulatory cell activity across olfactory structures in *Drosophila melanogaster*. 522177.
91. Zhang X, Gaudry QJE. (2016). Functional integration of a serotonergic neuron in the *Drosophila* antennal lobe. 5:e16836.
92. Zhao XC, Berg BGJCs. (2009). Morphological and physiological characteristics of the serotonin-immunoreactive neuron in the antennal lobe of the male oriental tobacco budworm, *Helicoverpa assulta*. 34(5):363-372.
93. Berck ME, Khandelwal A, Claus L, Hernandez-Nunez L, Si G, Tabone CJ, et al. (2016). The wiring diagram of a glomerular olfactory system. 5:e14859.
94. Coates KE, Majot AT, Zhang X, Michael CT, Spitzer SL, Gaudry Q, et al. (2017). Identified serotonergic modulatory neurons have heterogeneous synaptic connectivity within the olfactory system of *Drosophila*. 37(31):7318-7331.
95. Sun X, Tolbert L, Hildebrand JJJoCN. (1993). Ramification pattern and ultrastructural characteristics of the serotonin-immunoreactive neuron in the antennal lobe of the moth *Manduca sexta*: a laser scanning confocal and electron microscopic study. 338(1):5-16.
96. Ogawa SK, Cohen JY, Hwang D, Uchida N, Watabe-Uchida MJCr. (2014). Organization of monosynaptic inputs to the serotonin and dopamine neuromodulatory systems. 8(4):1105-1118.
97. Dorocic IP, Fürth D, Xuan Y, Johansson Y, Pozzi L, Silberberg G, et al. (2014). A whole-brain atlas of inputs to serotonergic neurons of the dorsal and median raphe nuclei. 83(3):663-678.

98. Fornal CA, Metzler CW, Marrosu F, Ribiero-do-Valle LE, Jacobs BLJBr. (1996). A subgroup of dorsal raphe serotonergic neurons in the cat is strongly activated during oral-buccal movements. 716(1-2):123-133.
99. Gao K, Mason PJJoN. (2000). Serotonergic raphe magnus cells that respond to noxious tail heat are not ON or OFF cells. 84(4):1719-1725.
100. Moriya S, Yamashita A, Nishi R, Ikoma Y, Yamanaka A, Kuwaki TJr. (2019). Acute nociceptive stimuli rapidly induce the activity of serotonin and noradrenalin neurons in the brain stem of awake mice. 7:1-9.
101. Rasmussen K, Heym J, Jacobs BLJEn. (1984). Activity of serotonin-containing neurons in nucleus centralis superior of freely moving cats. 83(2):302-317.
102. Rasmussen K, Strecker RE, Jacobs BLJBr. (1986). Single unit response of noradrenergic, serotonergic and dopaminergic neurons in freely moving cats to simple sensory stimuli. 369(1-2):336-340.
103. Ranade SP, Mainen ZFJJon. (2009). Transient firing of dorsal raphe neurons encodes diverse and specific sensory, motor, and reward events. 102(5):3026-3037.
104. Ren J, Friedmann D, Xiong J, Liu CD, Ferguson BR, Weerakkody T, et al. (2018). Anatomically defined and functionally distinct dorsal raphe serotonin sub-systems. 175(2):472-487. e20.
105. Waterhouse BD, Devilbiss D, Seiple S, Markowitz RJBr. (2004). Sensorimotor-related discharge of simultaneously recorded, single neurons in the dorsal raphe nucleus of the awake, unrestrained rat. 1000(1-2):183-191.
106. Lizbinski KM, Dacks AMJFinc. (2018). Intrinsic and extrinsic neuromodulation of olfactory processing. 11:424.
107. Haider B, Häusser M, Carandini MJN. (2013). Inhibition dominates sensory responses in the awake cortex. 493(7430):97-100.
108. Mukaida K, Shichino T, Koyanagi S, Himukashi S, Fukuda KJA, Analgesia. (2007). Activity of the serotonergic system during isoflurane anesthesia. 104(4):836-839.
109. Portas CM, Krakow K, Allen P, Josephs O, Armony JL, Frith CDJN. (2000). Auditory processing across the sleep-wake cycle: simultaneous EEG and fMRI monitoring in humans. 28(3):991-999.
110. Azimi Z, Barzan R, Spoida K, Surdin T, Wollenweber P, Mark MD, et al. (2020). Separable gain control of ongoing and evoked activity in the visual cortex by serotonergic input. 9:e53552.
111. Berkes P, Orbán G, Lengyel M, Fiser JJS. (2011). Spontaneous cortical activity reveals hallmarks of an optimal internal model of the environment. 331(6013):83-87.
112. Fiser J, Berkes P, Orbán G, Lengyel MJTics. (2010). Statistically optimal perception and learning: from behavior to neural representations. 14(3):119-130.
113. Lottem E, Lörincz ML, Mainen ZFJJoN. (2016). Optogenetic activation of dorsal raphe serotonin neurons rapidly inhibits spontaneous but not odor-evoked activity in olfactory cortex. 36(1):7-18.
114. Hurley L, Devilbiss D, Waterhouse BJCoin. (2004). A matter of focus: monoaminergic modulation of stimulus coding in mammalian sensory networks. 14(4):488-495.
115. Mooney R, Huang X, Shi M-Y, Bennett-Clarke C, Rhoades RJPibr. (1996). Serotonin modulates retinotectal and corticotectal convergence in the superior colliculus. 112:57-69.
116. Petzold GC, Hagiwara A, Murthy VNJNn. (2009). Serotonergic modulation of odor input to the mammalian olfactory bulb. 12(6):784-791.
117. Salinas E, Thier PJN. (2000). Gain modulation: a major computational principle of the central nervous system. 27(1):15-21.
118. Cohen JY, Amoroso MW, Uchida NJE. (2015). Serotonergic neurons signal reward and punishment on multiple timescales. 4:e06346.
119. Jacob SN, Nienborg HJFinc. (2018). Monoaminergic neuromodulation of sensory processing. 12:51.
120. Costa VD, Kakalios LC, Averbeck BJBn. (2016). Blocking serotonin but not dopamine reuptake alters neural processing during perceptual decision making. 130(5):461.
121. Chen C, Regehr WGJJoN. (2003). Presynaptic modulation of the retinogeniculate synapse. 23(8):3130-3135.
122. Seeburg DP, Liu X, Chen CJJoN. (2004). Frequency-dependent modulation of retinogeniculate transmission by serotonin. 24(48):10950-10962.
123. Pickard GE, Smith BN, Belenky M, Rea MA, Dudek FE, Sollars PJJoN. (1999). 5-HT1B receptor-mediated presynaptic inhibition of retinal input to the suprachiasmatic nucleus. 19(10):4034-4045.
124. Smith BN, Sollars PJ, Dudek FE, Pickard GEJJobr. (2001). Serotonergic modulation of retinal input to the mouse suprachiasmatic nucleus mediated by 5-HT1B and 5-HT7 receptors. 16(1):25-38.
125. Fitzgerald KK, Sanes DHJJon. (1999). Serotonergic modulation of synapses in the developing gerbil lateral superior olive. 81(6):2743-2752.
126. Filosa A, Barker AJ, Dal Maschio M, Baier HJN. (2016). Feeding state modulates behavioral choice and processing of prey stimuli in the zebrafish tectum. 90(3):596-608.
127. Yoshida M, Sasa M, Takaori SJBr. (1984). Serotonin-mediated inhibition from dorsal raphe nucleus of neurons in dorsal lateral geniculate and thalamic reticular nuclei. 290(1):95-105.
128. Reggiani JD, Jiang Q, Barbini M, Lutas A, Liang L, Fernando J, et al. (2023). Brainstem serotonin neurons selectively gate retinal information flow to thalamus. 111(5):711-26. e11.
129. Mooney RD, Shi MY, Rhoades RWJJoN. (1994). Modulation of retinotectal transmission by presynaptic 5-HT1B receptors in the superior colliculus of the adult hamster. 72(1):3-13.
130. Laurent A, Goaillard J-M, Cases O, Lebrand C, Gaspar P, Ropert NJJoN. (2002). Activity-dependent presynaptic effect of serotonin 1B receptors on the somatosensory thalamocortical transmission in neonatal mice. 22(3):886-900.
131. Araneda R, Andrade RJN. (1991). 5-Hydroxytryptamine₂ and 5-hydroxytryptamine_{1A} receptors mediate opposing responses on membrane excitability in rat association cortex. 40(2):399-412.
132. Tanaka E, North RAJJon. (1993). Actions of 5-hydroxytryptamine on neurons of the rat cingulate cortex. 69(5):1749-1757.
133. Sheldon PW, Aghajanian GKJS. (1991). Excitatory responses to serotonin (5-HT) in neurons of the rat piriform cortex:

- Evidence for mediation by 5-HT_{1C} receptors in pyramidal cells and 5-HT₂ receptors in interneurons. 9(3):208-218.
134. Marek GJ, Aghajanian GKJJoP, Therapeutics E. (1996). LSD and the phenethylamine hallucinogen DOI are potent partial agonists at 5-HT_{2A} receptors on interneurons in rat piriform cortex. 278(3):1373-1382.
 135. Zhou F-M, Hablitz JJJJon. (1999). Activation of serotonin receptors modulates synaptic transmission in rat cerebral cortex. 82(6):2989-2999.
 136. Celada P, Puig MV, Casanovas JM, Guillazo G, Artigas FJJJoN. (2001). Control of dorsal raphe serotonergic neurons by the medial prefrontal cortex: involvement of serotonin-1A, GABAA, and glutamate receptors. 21(24):9917-9929.
 137. Brigman JL, Mathur P, Harvey-White J, Izquierdo A, Saksida LM, Bussey TJ, et al. (2010). Pharmacological or genetic inactivation of the serotonin transporter improves reversal learning in mice. 20(8):1955-1963.
 138. Zhou FC, Lesch K-P, Murphy DLJBr. (2002). Serotonin uptake into dopamine neurons via dopamine transporters: a compensatory alternative. 942(1-2):109-119.
 139. Goel A, Cantu DA, Guilfoyle J, Chaudhari GR, Newadkar A, Todisco B, et al. (2018). Impaired perceptual learning in a mouse model of Fragile X syndrome is mediated by parvalbumin neuron dysfunction and is reversible. 21(10):1404-1411.
 140. Kissinger ST, Wu Q, Quinn CJ, Anderson AK, Pak A, Chubykin AAJCr. (2020). Visual experience-dependent oscillations and underlying circuit connectivity changes are impaired in Fmr1 KO mice. 31(1).
 141. Xiang Z, Prince DAJJJoN. (2003). Heterogeneous actions of serotonin on interneurons in rat visual cortex. 89(3):1278-1287.
 142. Patel AM, Kawaguchi K, Seillier L, Nienborg HJEJoN. (2023). Serotonergic modulation of local network processing in V1 mirrors previously reported signatures of local network modulation by spatial attention. 57(8):1368-1382.
 143. Azizi S, Waterhouse B, Burne R, Woodward DJIOVS. (1983). Modulatory actions of norepinephrine and serotonin on responses of simple and complex cells in rat visual cortex. 24(228):407-416.
 144. Ferron A, Descarries L, Reader TAJBR. (1982). Altered neuronal responsiveness to biogenic amines in rat cerebral cortex after serotonin denervation or depletion. 231(1):93-108.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

Submit Manuscript

DOI:10.31579/2578-8868/336

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://auctoresonline.org/journals/neuroscience-and-neurological-surgery>