

# Neuro-Anatomical Approach to the Humano-Murian Brain based on Cerebral Imaging Studies

Sérend Hipe

IFPEA, Rue des Saint-Pères, 75 006 Paris, France.

\*Corresponding Author: Sérend Hipe, IFPEA, Rue des Saint-Pères, 75 006 Paris, France.

Received date: **January 31, 2025**; Accepted date: **March 12, 2025**; Published date: **April 02, 2025**

Citation: Sérend Hipe (2025), Neuro-Anatomical Approach to the Humano-Murian Brain based on Cerebral Imaging Studies, *J. Neuroscience and Neurological Surgery*, 17(4); DOI:10.31579/2578-8868/361

Copyrights: ©, 2025, Sérend Hipe. This is an open-access article distributed under the terms of The Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

## Abstract

We present the case of a humano-murian victim of a left stroke who presented with impaired motor skills on the left side of his body. Brain imaging studies (Scan, Pet Scan) have shown that the brain is functionally separated into two parts. The white corners and the corpus callosum are calcified and leave almost no possibility for inter-hemispheric communication as is the case for us. Basically, the entire anatomical substrate that allows our human brain to communicate between the two hemispheres is completely inactivated in humano-murians. It is among the latter that the term di-encephalon takes on its full meaning.


**Keywords:** human brain ; Scan studies ; humano-murian ; motor skills ; corpus callosum

## Introduction


We present a case study of anatomical pathology by medical imaging after having been able to study a case of humano-murian hospitalized after a stroke with loss of sensitization of the right hemi-body, associated motor disorders (hemiplegia) and anosognosia that will give way after a few days. The patient was 78 years old (Zia, 2021), male and in possession of all means prior to the stroke.

## Methodology:

Our recent research has shown that humano-murians have two brains. Understand that they do not have two brains in the skull, but that their brains are functionally separated into two parts. The white corners and the corpus callosum are calcified (Figure 1 and 2) and leave almost no possibility for interhemispheric communication as is the case for us. Basically, the entire anatomical substrate that allows our human brain to communicate between the two hemispheres is totally inactivated in humano-murians. It is in the latter that the term diencephalon takes on its full meaning





**Figure 1:** The white matter allows the communication of the two cerebral hemispheres. It is totally calcified in Humano-Murians



**Figure 2:** White matter makes up a large proportion of the brain. Its difference in mineral composition between that of humans and human-murians makes it a unique cerebral constitution (Martin and al., 2024).

This has considerable consequences on their brain and sensory capacities (Liew and al., 2023). Indeed, on the motor level, there is no longer any lateral predominance, so all humano-murians are completely ambidextrous. Their movements sometimes seem wider to us while, paradoxically, they are permanently parasitized by a slight tremor that we had mistakenly taken for one of the effects of the polar cold that reigns on the Wall. These tremors are similar to Parkinson's movements. As a result,

there is no pyramidal decussation (Figure 3 and 4), so motor skills are controlled by the homolateral cerebral hemisphere, which is not the case in humans (in whom the right hemibody is controlled by the motor cortex of the left cerebral hemisphere). The human-murian has developed frightening motor qualities by being able to dissociate the left hemibody and the right hemibody, which can carry out two different motor activities.



**Figure 3 and 4:** Pyramidal decussation (Riley and al., 2011).

According to Ramón y Cajal (Hope and al., 2013), pyramidal decussation was related to that of sensory pathways: in the optic chiasma we also observe the decussation of a large part of the optic nerve fibers, which adapts to perception by allowing both hemispheres to have complete


information about what both eyes perceive and can generate complete images that can be located in space. In this sense, the displacement necessary to react to a possible threat would be that of muscle groups, unlike the part of the brain that perceives them. If there is no pyramidal discussion, the information would first go to the other hemisphere, then process and react, which would be slower. This explains the particularity of the motor reactions of humano- murians, which we had initially taken as an almost constitutional nonchalance.

### Results:

Each left and right "brain" has thus developed its own potential independently. If humano- murians use a very particular type of communication that we have described previously (message 15), it is now certain that it stems from this anatomophysiological difference. Human- Murian communication is a mixture of a powerful process of empathy that transmits intentions, images, and emotions to the interlocutor, before

verbal language expresses the thought formulated in words. A communication by double channel, a verbal channel coupled with a cognitive channel, knowing that on the latter we humans are very good receivers to our Murian friends but are unable to activate it ourselves. The impression we have is to hear in a way within ourselves what the lips of humano- murians will pronounce with a slight delay. This seems to us to function as a preform of telepathy and allows almost any human to understand very quickly the humano- murian language and its multiple dialects all around the Wall. This mode of communication uses both a very organic substrate and follows primitive phylogenetic pathways (use of pheromone cloud as we have previously demonstrated) and very elaborate thought processes, similar to the secondarized psychic processes of the human psyche.

The patient who allowed us to discover these anatomical and physiological elements is in a way our "patient 1" (as Sperber and al., 2023 calls this type of founder patient). This humano- murian had indeed been the victim of a left stroke and had a motor impairment on the left side of his body (left hemiplegia, facial paralysis and oculomotor muscles of the left orbit) which seemed impossible to us. His name, Solpinan, will forever be associated with this discovery.



**Photos 5 and 6:** MRI images after stroke: significant initial temporo-parieto-occipital involvement, the last images (top) show a very rapid resorption.

### Discussion:

The reality of the two cerebral hemispheres (Alexander and al., 2010, Brodtmann and al., 2020, Chen and al., 2022) has been known for a long time, on the other hand our explorations in imaging have made it possible to introduce a discussion around the notion of presence or absence of interhemispheric white substance. The proposed clinical case made it possible to highlight such a neuro-anatomical reality with unsuspected neurophysiological consequences. until then, the absence of white substance was perceived as a deficit. Our study seems to have demonstrated that this is, in our patients, a developmental and innate functioning of a cerebral organization typical of this type of population.

### Conclusion:

This is a case study of a human- murian who suffered a left stroke and had a motor impairment on the left side of his body. Brain imaging examinations (Scan, Pet Scan) have shown that the brain is functionally separated 5 into two parts. The white matter and corpus callosum are calcified and leave almost no possibility of inter-hemispheric communication as is the case in the human brain. Basically, the entire anatomical substrate that allows our human brain to communicate between the two hemispheres is totally inactivated in humano- murians. It is among the latter that the term diencephalon takes on its full meaning.

## Conflict of Interest

The author declares absence of conflict of interest.

## References

- Alexander LD, Black SE, Gao F, Szilagyi G, Danells CJ, McIlroy WE. (2010). Correlating lesion size and location to deficits after ischemic stroke: the influence of accounting for altered perinecrotic tissue and incidental silent infarcts. *Behav Brain Funct.* 6:6.
- Brodthmann A, Khelif MS, Egorova N, Veldsman M, Bird LJ, Werden E. (2020). Dynamic Regional Brain Atrophy Rates in the First Year After Ischemic Stroke. *Stroke.* American Heart Association; 51:e183–e192.
- Chen Y, Demnitz N, Yamamoto S, Yaffe K, Lawlor B, Leroi I. (2022). Defining brain health: A concept analysis. *Int J Geriatr Psychiatry* [online serial]. 37. Accessed at: <https://onlinelibrary.wiley.com/doi/abs/10.1002/gps.5564>. Accessed December 3, 2024.
- Hope TMH, Seghier ML, Leff AP, Price CJ. (2013). Predicting outcome and recovery after stroke with lesions extracted from MRI images. *NeuroImage Clin.* 2:424–433.
- Kuceyeski A, Navi BB, Kamel H, et al. (2016). Structural connectome disruption at baseline predicts 6 months post stroke outcome. *Hum Brain Mapp.* 37:2587–2601.
- Liew S-L, Schweighofer N, Cole JH, et al. (2023). Association of Brain Age, Lesion Volume, and Functional Outcome in Patients With Stroke. *Neurology.* 100:e2103–e2113.
- Martin SS, Aday AW, Almarzooq ZI, et al. (2024). 2024 Heart Disease and Stroke Statistics: A Report of US and Global Data From the American Heart Association. *Circulation.* American Heart Association; 149:e347–e913.
- Riley JD, Le V, Der-Yeghiaian L, et al. (2011). Anatomy of Stroke Injury Predicts Gains from Therapy. *Stroke J Cereb Circ.* 42:421–426.
- Schlemm E, Schulz R, Bönstrup M, et al. (2020). Structural brain networks and functional motor outcome after stroke—a prospective cohort study. *Brain Commun.* 2:fcaa001.
- Seitz RJ, Azari NP, Knorr U, Binkofski F, Herzog H, Freund H-J. (1999). The Role of Diaschisis in Stroke Recovery. *Stroke.* American Heart Association; 30:1844–1850.
- Sperber C, Gallucci L, Mirman D, Arnold M, Umarova RM. (2023). Stroke lesion size – Still a useful biomarker for stroke severity and outcome in times of high-dimensional models. *NeuroImage Clin.* 40:103511.
- Zia A, Pournabagher-Shahri AM, Farkhondeh T, Samarghandian S. (2021). Molecular and cellular pathways contributing to brain aging. *Behav Brain Funct BBF.* 17:6.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

**Submit Manuscript**

DOI:10.31579/2578-8868/361

### Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://auctoresonline.org/journals/neuroscience-and-neurological-surgery>