

Giant Gastric Trichobezoar: Subtotal Gastrectomy

Eduardo DomínguezAdame ^{1,3*}, Cristina Rubio Castellanos ¹, Manuel Rodríguez Téllez ^{2,3}

¹Departments of General and Digestive Surgery.

²Hepatogastroenterology.

³Virgen Macarena University Hospital. University of Seville.

***Corresponding Author:** Eduardo DomínguezAdame, Departments of General and Digestive Surgery Virgen Macarena University Hospital. University of Seville.

Received Date: August 07, 2025 | **Accepted Date:** September 04, 2025 | **Published Date:** September 23, 2025

Citation: Eduardo DomínguezAdame, Cristina R. Castellanos, Manuel R. Téllez, (2025), Giant Gastric Trichobezoar: Subtotal Gastrectomy, *International Journal of Clinical Case Reports and Reviews*, 30(2); DOI:10.31579/2690-4861/824

Copyright: © 2025, Eduardo DomínguezAdame. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract:

22-year-old patient (height 184 cm, weight 53 kg, Body Mass Index 15.65 kg/m²) with a previous history of trichophagia (Rapunzel Syndrome) in adolescence. For 1 year, he has had intermittent, stabbing epigastric discomfort of variable duration, unrelated to food intake and accompanied by hyporexia, nausea without vomiting, premature satiety and unquantified weight loss. He has a painless palpable tumor in the epigastrium

Key words: giant gastric trichobezoar; hyporexia

Introduction

22-year-old patient (height 184 cm, weight 53 kg, Body Mass Index 15.65 kg/m²) with a previous history of trichophagia (Rapunzel Syndrome) in adolescence. For 1 year, he has had intermittent, stabbing epigastric

discomfort of variable duration, unrelated to food intake and accompanied by hyporexia, nausea without vomiting, premature satiety and unquantified weight loss. He has a painless palpable tumor in the epigastrium. Endoscopy (Figure 1):




Figure 1: Endoscopic image of a Giant Gastric Trichobezoar. (A) Trichobezoar imprint in the cardia. (B) Stony, calcified gastric trichobezoar that prevents progression of the endoscope in the gastric cavity.

Megadilated stomach with a giant stony, calcified trichobezoar, not amenable to endoscopic treatment. Abdomino-pelvic CT scan (Figure 2)




Figure 2: CT scan Giant gastric trichobezoar. (A) Coronal image. (B) Axial image. (C) Sagittal image.

Stomach enlargement, reaching 250 mm longitudinally, extending over the void and right iliac fossa, not ruling out that they correspond to food remains/foreign body. She was referred for surgery (Figure 3)




Figure 3: Subtotal gastrectomy for Giant Gastric Trichobezoar. (A) Stomach almost completely occupied. (B) Stony, calcified and impacted gastric trichobezoar.

Where a megastomach was found due to a giant, calcified, impacted stony trichobezoar. A subtotal gastrectomy was performed with a Roux-en-Y loop gastrojejunal anastomosis (Billroth III technique). The patient was discharged from the hospital on the fourth day without incident.

Discussion

Gastric trichobezoars can become large and extend into the small intestine (Rapunzel syndrome). Their diagnosis is completed with imaging studies such as ultrasound, CT scans, and endoscopies. Treatment is based on the removal of the trichobezoar, either endoscopically or surgically. Sometimes, due to its size, impaction, and time of evolution, it will be necessary to perform an excisional surgery (gastrectomy) due to the gastric atony caused by the trichobezoar [1,2,3].

References

1. Paschos KA, Chatzigeorgiadis A.(2019). Pathophysiological and clinical aspects of the diagnosis and treatment of bezoars. *Ann Gastroenterol*;32(3):224-232.
2. Khalifa MB, Ghannouchi M, Nacef K.(2022). Trichobezoar: a case report of a double gastric and ilial localization revealed by an occlusion. *Int J Surg Case Rep*;91:106782.
3. Luisa Paola Garzón Hernández, Isabel Mora Oliver, Ernesto Muñoz Sornosa, Elena Martí Cuñat.(2023). Duodenal diastatic perforation due to double gastric and jejunal trichobezoar in a patient with Rapunzel syndrome. *Rev Esp Enferm Dig*. 2023;115(4):222-223.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

Submit Manuscript

DOI: [10.31579/2690-4861/824](https://doi.org/10.31579/2690-4861/824)

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://auctoresonline.org/journals/international-journal-of-clinical-case-reports-and-reviews>